

# On path-dependent SDEs involving distributional drifts

Alberto Ohashi<sup>a</sup>, Francesco Russo<sup>b,\*</sup>, Alan Teixeira<sup>b</sup>

<sup>a</sup>*Departamento de Matemática, Universidade de Brasília, 70910-900, Brasília, Brazil*

<sup>b</sup>*ENSTA Paris, Institut Polytechnique de Paris, Unité de Mathématiques appliquées, 828, boulevard des Maréchaux, F-91120 Palaiseau, France*

[amfohashi@gmail.com](mailto:amfohashi@gmail.com) (A. Ohashi), [francesco.russo@ensta-paris.fr](mailto:francesco.russo@ensta-paris.fr) (F. Russo),  
[alan.teixeira@ensta-paris.fr](mailto:alan.teixeira@ensta-paris.fr) (A. Teixeira)

Received: 25 April 2021, Revised: 9 December 2021, Accepted: 9 December 2021,  
Published online: 3 January 2022

**Abstract** The paper presents the study on the existence and uniqueness (strong and in law) of a class of non-Markovian SDEs whose drift contains the derivative in the sense of distributions of a continuous function.

**Keywords** SDEs with distributional drift, path-dependent stochastic differential equations  
**2010 MSC** 60G99, 60H10, 60H15

## 1 Introduction

This paper discusses in detail a framework of one-dimensional stochastic differential equations (henceforth abbreviated by SDEs) with distributional drift and possible path-dependency. To our best knowledge, this is the first paper which approaches a class of non-Markovian SDEs with distributional drifts.

The main objective of this paper is to analyze the solution (existence and uniqueness) of the martingale problem associated with SDEs of the type

$$dX_t = \sigma(X_t)dW_t + b'(X_t)dt + \Gamma(t, X^t)dt, \quad X_0 \stackrel{d}{=} \delta_{x_0}, \quad (1.1)$$

---

\*Corresponding author.

where  $b, \sigma : \mathbb{R} \rightarrow \mathbb{R}$  are continuous functions,  $\sigma > 0$ ,  $x_0 \in \mathbb{R}$  and  $W$  is a standard Brownian motion. The assumptions on  $b$ , which will be formulated later, imply that  $b'$  is a Schwartz distribution. Concerning the path-dependent component of the drift, we consider a locally bounded functional

$$\Gamma : \Lambda \rightarrow \mathbb{R}, \quad (1.2)$$

where

$$\Lambda := \{(s, \eta) \in [0, T] \times C([0, T]); \eta = \eta^s\}$$

and

$$\eta^s(t) := \begin{cases} \eta(t), & \text{if } t \leq s, \\ \eta(s), & \text{if } t > s, \end{cases}$$

and  $C([0, T])$  is the space of real-valued continuous functions on  $[0, T]$ , for a given terminal time  $0 < T < \infty$ . By convention, we extend  $\Gamma$  from  $\Lambda$  to  $[0, T] \times C([0, T])$  by setting (in nonanticipating way)

$$\Gamma(t, \eta) := \Gamma(t, \eta^t), t \in [0, T], \eta \in C([0, T]). \quad (1.3)$$

Setting  $\sigma = 1$ ,  $b = B$ , where  $B$  is a two-sided real-valued Brownian motion which is independent from  $W$ , equation (1.1) reads

$$dX_t = -\frac{1}{2}\dot{B}(X_t)dt + \Gamma(t, X^t)dt + dW_t. \quad (1.4)$$

When  $\Gamma = 0$ , equation (1.4) constitutes the so-called Brox diffusion, see, e.g., [4, 15, 25] and other references therein. This is a celebrated *random environment* model. This paper includes the study of (1.4), where  $\Gamma$  is a bounded path-dependent functional, which appears to be a non-Markovian variant of Brox diffusion.

Path-dependent SDEs were investigated under several aspects. Under standard Lipschitz regularity conditions on the coefficients, it is known (see, e.g., Theorem 11.2 [20, chapter V]) that strong existence and uniqueness hold. In the case the path-dependence takes the form of delayed stochastic equations, one-sided Lipschitz condition ensures strong existence and uniqueness, see, e.g., [24, 17]. Beyond Lipschitz regularity on the coefficients of the SDE, [14] shows uniqueness in law under structural conditions on an underlying approximating Markov process, where local-time and running maximum dependence are considered. The existence in law for infinite-dimensional SDEs with additive noise on the configuration space with path-dependent drift functionals with sublinear growth is studied in [8]. In all those contributions the drift is a nonanticipative functional. Beyond Brownian motion based driving noises, [6] establishes existence of solutions for some path-dependent differential equation driven by a Hölder process.

The Markovian case ( $\Gamma = 0$ ) with distributional drift has been intensively studied over the years. Diffusions in the generalized sense were first considered in the case when the solution is still a semimartingale, beginning with [19]. Later on, many authors considered special cases of SDEs with generalized coefficients. It is difficult to quote them all; in the case when the drift  $b'$  is a measure and the solutions are semimartingales we refer the reader to [3, 9, 21]. We also recall that [10] considered

even special cases of nonsemimartingales solving stochastic differential equations with generalized drift.

In [12] and [13], the authors studied time-independent one-dimensional SDEs of the form

$$dX_t = \sigma(X_t)dW_t + b'(X_t)dt, \quad t \in [0, T], \quad (1.5)$$

whose solutions are possibly nonsemimartingale processes, where  $\sigma$  is a strictly positive continuous function and  $b'$  is the derivative of a real-valued continuous function. They presented well-posedness of the martingale problem, Itô's formula under weak conditions, semimartingale characterization and the Lyons–Zheng decomposition. The only supplementary assumption was the existence of the function

$$\Sigma(x) := 2 \int_0^x \frac{b'}{\sigma^2}(y)dy, \quad x \in \mathbb{R}, \quad (1.6)$$

considered as a suitable limit via regularizations. Those authors considered solutions in law. The SDE (1.5) was also investigated in [1], where the authors provided a well-stated framework when  $\sigma$  and  $b$  are  $\gamma$ -Hölder continuous,  $\gamma > \frac{1}{2}$ . In [22], the authors have also shown that in some cases strong solutions exist and pathwise uniqueness holds. More recently, in the time-dependent framework (but still one-dimensional), a significant contribution was done in [7]. As far as the multidimensional case is concerned, some important steps were done in [11] and more recently in [5], when the diffusion matrix is the identity and  $b$  is a time-dependent drift in some proper negative Sobolev space. We also refer to [2], where the authors have focused on (1.1) in the case of a time-independent drift  $b$  which is a measure of Kato class.

Let us come back to the objective of the present paper in which the path-dependent drift contains the derivative in the sense of distributions of a continuous function  $b$ . We remark that in case  $b'$  is a bounded measurable function and  $\Gamma$  is a bounded path-dependent functional, then the problem can be easily treated by applying Girsanov's theorem, see, e.g., [16, Proposition 3.6 and 3.10, chapter 5]. Here, the combination of a Schwartz distribution  $b'$  with a path-dependent functional  $\Gamma$  requires a new set of ideas. Equation (1.1) will be interpreted as a martingale problem with respect to an operator

$$\mathcal{L}f := Lf + \Gamma f',$$

see (3.3), where  $L$  is the Markovian generator

$$Lf = \frac{\sigma^2}{2} f'' + b' f', \quad (1.7)$$

where we stress that  $b'$  is the derivative of some continuous function  $b$ . If we define  $\Sigma$  as in (1.6), then the operator  $L$  can be written as

$$Lf = (e^\Sigma f')' \frac{e^{-\Sigma} \sigma^2}{2}, \quad (1.8)$$

see [12]. We define a notion of martingale problem related to  $\mathcal{L}$  (see Definition 3.3) and a notion of strong martingale problem related to  $\mathcal{D}_{\mathcal{L}}$  and a given Brownian motion  $W$  (see Definition 3.4). That definition has to be compared with the notion of strong

existence and pathwise uniqueness of an SDE. In this article, the notion of martingale problem extends the usual one by replacing the space  $C^2$  of twice continuously differentiable real-valued functions with a more suitable set  $\mathcal{D}_{\mathcal{L}}$ . In the Markovian case, the notion of *strong martingale problem* was introduced in [22]. As mentioned earlier, we will concentrate on the case when  $b$  is continuous, the case of special discontinuous functions is investigated in [18].

The strategy of this paper consists in eliminating the distributional drift of the so-called Zvonkin's transform, see [27]. In this direction, we transform the equation via an  $L$ -harmonic function  $h$  which exists under the assumption that the function (1.6) is well-defined. The case with  $\Gamma = 0$  was already implemented in [12] and [13], where the drift in the transformed SDE was null. In our non-Markovian context, the transformed equation is essentially a path-dependent SDE with measurable coefficients. Under some linear growth conditions (see Assumption 4.16), Theorem 4.23 illustrates the existence of the martingale problem related to (1.1). Proposition 4.24 states uniqueness under more restrictive conditions. Indeed, consider the example when  $\sigma = 1$  and  $b'$  is the derivative (in the sense of distributions) of a bounded continuous function  $b$  and  $\Gamma$  is a bounded measurable functional. In this case,  $h = \int_0^\cdot e^{-2b(y)} dy$ . Then the study of the well-posedness of the martingale problem is equivalent to the well-posedness of the path-dependent SDE

$$Y = Y_0 + \int_0^\cdot \exp(-2b(h^{-1}(Y_s))) dW_s + \int_0^\cdot \Gamma(s, h^{-1}(Y_s)) \exp(-2b(h^{-1}(Y_s))) ds. \quad (1.9)$$

Existence and uniqueness for (1.9) can be established via Girsanov's theorem.

Moreover, Corollary 4.31 establishes well-posedness for the strong martingale problem associated to (1.1). This holds under suitable Lipschitz regularity conditions on the functional  $\tilde{\Gamma}$ , which is related to  $\Gamma$  via (4.14), and a specific assumption on the function  $\sigma_0$  defined in Remark 4.9. We suppose that  $\sigma_0$  is bounded, uniformly elliptic and it fulfills a Yamada–Watanabe type condition. One typical example is given when  $\sigma_0$  is a  $\gamma$ -Hölder continuous function for  $\frac{1}{2} \leq \gamma \leq 1$ .

Several results of the present paper can be partially extended to the multidimensional case, by using the techniques developed in the Markovian case in [11]. In this direction, the harmonic function  $h$  should be replaced by the “mild” solution  $\phi$  of the parabolic Kolmogorov equation

$$\begin{cases} \partial_t \phi + \frac{1}{2} \Delta \phi + b' \nabla \phi & = \lambda(\phi - \text{id}), \\ \phi(T, \cdot) & = \text{id}, \end{cases} \quad (1.10)$$

see Section 3.2 of [11].

## 2 Notations and preliminaries

### 2.1 General notations

Let  $I$  be an interval of  $\mathbb{R}$ .  $C^k(I)$  is the space of real functions defined on  $I$  having continuous derivatives till order  $k$ . Such space is endowed with the topology of the uniform convergence on compact sets for the functions and all derivatives. Generally,

$I = \mathbb{R}$  or  $[0, T]$  for some fixed positive real  $T$ . The space of continuous functions on  $I$  will be denoted by  $C(I)$ . Often, if there is no ambiguity,  $C^k(\mathbb{R})$  will be simply indicated by  $C^k$ . Given an a.e. bounded real function  $f$ ,  $|f|_\infty$  will denote the essential supremum.

We recall some notions from [12]. For us, all filtrations  $\mathfrak{F}$  fulfill the usual conditions. When no filtration is specified, we mean the canonical filtration of the underlying process. Otherwise the canonical filtration associated with a process  $X$  is denoted by  $\mathfrak{F}^X$ . An  $\mathfrak{F}$ -Dirichlet process  $X$  is the sum of an  $\mathfrak{F}$ -local martingale  $M^X$  with an  $\mathfrak{F}$ -adapted zero quadratic variation process  $A^X$ . We will fix by convention that  $A_0^X = 0$  so that the decomposition is unique. A sequence  $(X^n)$  of continuous processes indexed by  $[0, T]$  is said to converge u.c.p. to some process  $X$  whenever  $\sup_{t \in [0, T]} |X_t^n - X_t|$  converges to zero in probability. Finally the notion of covariation between two general càdlàg processes (whenever it exists) is denoted by  $[X, Y]$  and we set  $[X] = [X, X]$ , see, e.g., [23]. If  $[X]$  exists,  $X$  is called a finite quadratic variation process.

**Remark 2.1.** (1) An  $\mathfrak{F}$ -continuous semimartingale  $Y$  is always an  $\mathfrak{F}$ -Dirichlet process. The  $A^Y$  process coincides with the continuous bounded variation component. Moreover, the quadratic variation  $[Y]$  is the usual quadratic variation for semimartingales.

(2) Any  $\mathfrak{F}$ -Dirichlet process is a finite quadratic variation process and its quadratic variation is given by  $[X] = [M^X]$ .

(3) If  $f \in C^1(\mathbb{R})$  and  $X = M^X + A^X$  is an  $\mathfrak{F}$ -Dirichlet process, then  $Y = f(X)$  is again an  $\mathfrak{F}$ -Dirichlet process and  $[Y] = \int_0^\cdot f'(X)^2 d[M^X]$ .

### 3 Non-Markovian SDE: the function case

#### 3.1 General considerations

As in the case of Markovian SDEs, it is possible to formulate the notions of strong existence, pathwise uniqueness, existence and uniqueness in law for path-dependent SDEs of the type (1.1), see, e.g., Section A. Let us suppose for the moment that  $\sigma, b' : \mathbb{R} \rightarrow \mathbb{R}$  are Borel functions. We will consider solutions  $X$  of

$$\begin{cases} dX_t &= \sigma(X_t)dW_t + b'(X_t)dt + \Gamma(t, X^t)dt, \\ X_0 &= \xi, \end{cases} \quad (3.1)$$

for some initial condition  $\xi$ .

The previous equation will be denoted by  $E(\sigma, b', \Gamma; \nu)$  (where  $\nu$  is the law of  $\xi$ ), or simply by  $E(\sigma, b', \Gamma)$  if we omit the initial condition. For simplicity, the initial conditions will always be considered as deterministic. The functional  $\Gamma$  is nonanticipative in the sense of (1.3).

**Definition 3.1.** Let  $\nu$  be the Dirac probability measure on  $\mathbb{R}$  such that  $\nu = \delta_{x_0}$ ,  $x_0 \in \mathbb{R}$ . A stochastic process  $X$  is called a **solution** of  $E(\sigma, b', \Gamma; \nu)$  with respect to a probability  $\mathbb{P}$  if there is a Brownian motion  $W$  on some filtered probability space, such that  $X$  solves (3.1) and  $X_0 = x_0$ . We also say that the couple  $(X, \mathbb{P})$  solves  $E(\sigma, b', \Gamma)$  with initial condition distributed according to  $\nu$ .

Suppose  $\Gamma \equiv 0$ . A very well-known result in [26], Corollary 8.1.7, concerns the equivalence between martingale problems and solution in law of SDEs. Suppose for a moment that  $b'$  is a continuous function. According to [16, chapter 5], a process  $X$  and probability  $\mathbb{P}$  solve the classical martingale problem, if and only if,  $X$  is a solution of (1.1). The proof of that result can be easily adapted to the path-dependent case, i.e. when  $\Gamma \neq 0$ . This provides the statement below.

**Proposition 3.2.** *A couple  $(X, \mathbb{P})$  is a solution of  $E(\sigma, b', \Gamma)$ , if and only if, under  $\mathbb{P}$ ,*

$$f(X_t) - f(X_0) - \int_0^t Lf(X_s)ds - \int_0^t f'(X_s)\Gamma(s, X^s)ds \quad (3.2)$$

is a local martingale, where  $Lf = \frac{1}{2}\sigma^2 f'' + b' f'$ , for every  $f \in C^2$ .

### 3.2 Comments about the distributional case

When  $b'$  is a distribution, it is not obvious to introduce the notion of SDE, except in the case when  $L$  is close to the divergence form, i.e. when  $Lf = (\sigma^2 f')' + \beta f'$  and  $\beta$  is a Radon measure, see, e.g., Proposition 3.1 of [12]. For this reason, we replace the notion of solution in law with the notion of martingale problem. Suppose for a moment that  $L$  is a second order PDE operator with possibly generalized coefficients. In general, as it is shown in [12],  $C^2$  is not included in the natural domain of operator  $L$  and, similarly to [12], we will replace  $C^2$  with some set  $\mathcal{D}_L$ . Suppose that  $L : \mathcal{D}_L \subset C^1(\mathbb{R}) \rightarrow C(\mathbb{R})$ . Nevertheless  $\mathcal{D}_L$  is not the domain of  $L$  in the sense of the generator of a semigroup.

**Definition 3.3.** (1) We say that a continuous stochastic process  $X$  solves (with respect to a probability  $\mathbb{P}$  on some measurable space  $(\Omega, \mathcal{F})$ ) the martingale problem related to

$$\mathcal{L}f := Lf + \Gamma f', \quad (3.3)$$

with initial condition  $\nu = \delta_{x_0}$ ,  $x_0 \in \mathbb{R}$ , with respect to a domain  $\mathcal{D}_L$  if

$$M_t^f := f(X_t) - f(x_0) - \int_0^t Lf(X_s)ds - \int_0^t f'(X_s)\Gamma(s, X^s)ds \quad (3.4)$$

is a  $\mathbb{P}$ -local martingale for all  $f \in \mathcal{D}_L$ .

We will also say that the couple  $(X, \mathbb{P})$  is a solution of (or  $(X, \mathbb{P})$  solves) the martingale problem with respect to  $\mathcal{D}_L$ .

- (2) If a solution exists, we say that *existence* holds for the martingale problem above.
- (3) We say that *uniqueness holds* for the martingale problem above, if any two solutions (on some measurable space  $(\Omega, \mathcal{F})$ )  $(X^i, \mathbb{P}^i)$ ,  $i = 1, 2$ , have the same law.

We remark that in the classical literature of martingale problems, see [26], a solution is a probability on the path space  $C([0, T])$  and  $X$  is the canonical process. If  $(X, \mathbb{P})$  is a solution according to our notations, a solution in the classical framework

would be the probability law of  $X$  with respect to  $\mathbb{P}$ . We preserve our notations in conformity with [12, 13].

In the sequel, when the measurable space  $(\Omega, \mathcal{F})$  is self-explanatory it will be often omitted. As already observed in Proposition 3.2, the notion of martingale problem is (since the works of Stroock and Varadhan [26]) a concept related to solutions of SDEs in law. In the case when  $b'$  and  $\sigma$  are continuous functions (see [26]),  $\mathcal{D}_L$  corresponds to the space  $C^2(\mathbb{R})$ , in agreement with Remark 4.6 below.

Below we introduce the analogous notion of strong existence and pathwise uniqueness for our martingale problem.

**Definition 3.4.** (1) Let  $(\Omega, \mathcal{F}, \mathbb{P})$  be a probability space and let  $\mathfrak{F} = (\mathcal{F}_t)$  be the canonical filtration associated with a fixed Brownian motion  $W$ . Let  $x_0 \in \mathbb{R}$  be a constant. We say that a continuous  $\mathfrak{F}$ -adapted real-valued process  $X$  such that  $X_0 = x_0$  is a **solution to the strong martingale problem** (related to (3.3)), with respect to  $\mathcal{D}_L$  and  $W$  (with related filtered probability space), if

$$\begin{aligned} f(X_t) - f(x_0) - \int_0^t Lf(X_s)ds - \int_0^t f'(X_s)\Gamma(s, X^s)ds \\ = \int_0^t f'(X_s)\sigma(X_s)dW_s \end{aligned} \quad (3.5)$$

for all  $f \in \mathcal{D}_L$ .

- (2) We say that **strong existence** holds for the martingale problem related to (3.3) with respect to  $\mathcal{D}_L$ , if for every  $x_0 \in \mathbb{R}$ , given a filtered probability space  $(\Omega, \mathcal{F}, \mathbb{P}, \mathfrak{F})$ , where  $\mathfrak{F} = (\mathcal{F}_t)$  is the canonical filtration associated with a Brownian motion  $W$ , there is a process  $X$  solving the strong martingale problem (related to (3.3)) with respect to  $\mathcal{D}_L$  and  $W$  with  $X_0 = x_0$ .
- (3) We say that **pathwise uniqueness** holds for the martingale problem related to (3.3) with respect to  $\mathcal{D}_L$ , if given  $(\Omega, \mathcal{F}, \mathbb{P})$ , a Brownian motion  $W$  on it, two solutions  $X^i, i = 1, 2$ , to the strong martingale problem with respect to  $\mathcal{D}_L$  and  $W$ , and  $\mathbb{P}[X_0^1 = X_0^2] = 1$ ,  $X^1$  and  $X^2$  are indistinguishable.

## 4 The case when the drift is the derivative of a continuous function

### 4.1 The Markovian case

In this section we recall some basic notations and results from [12] but in a way that simplifies the presentation of our framework. We will also add some useful new elements. Let  $\sigma$  and  $b$  be functions in  $C(\mathbb{R})$  with  $\sigma > 0$ . In [12], in view of defining  $\mathcal{D}_L$  and  $L$  in the spirit of (1.7), the authors define the function

$$\Sigma(x) = 2 \lim_{n \rightarrow \infty} \int_0^x \frac{b'_n}{\sigma_n^2}(y)dy, \quad \forall x \in \mathbb{R}, \quad (4.1)$$

where the limit is intended to be in  $C(\mathbb{R})$ , i.e. uniformly on each compact. Here

$$\sigma_n^2 := \sigma^2 * \Phi_n, \quad b_n := b * \Phi_n, \quad (4.2)$$

where  $\Phi_n(x) := n\Phi(nx)$  and  $\Phi \in \mathcal{S}(\mathbb{R})$  (the Schwartz space), with  $\int \Phi(x) dx = 1$ . For concrete examples, one can take either  $\sigma^2$  or  $b$  to be locally bounded variation functions, see [12] for other examples and other details. Proposition 2.3, Lemma 2.6 and 2.9 of [12] allow us (equivalently) to define a subspace  $\mathcal{D}_L$  of  $C^1(\mathbb{R})$  on which the definition of  $Lf$  by (1.8) makes sense.

**Notation 4.1.** (1)  $\mathcal{D}_L$  is the subset of all  $f \in C^1(\mathbb{R})$  for which there exists  $\phi \in C^1$  such that  $f' = \exp(-\Sigma)\phi$ .

(2) If  $f \in \mathcal{D}_L$ , then we set

$$Lf = \phi' \exp(-\Sigma) \frac{\sigma^2}{2}, \quad (4.3)$$

where  $\phi$  is the function given in item (1) above.

(3) We denote by  $h : \mathbb{R} \rightarrow \mathbb{R}$  the function characterized by

$$h(0) = 0, \quad h' = e^{-\Sigma}. \quad (4.4)$$

In particular  $h$  is an  $L$ -harmonic function in the sense that  $Lh = 0$ , as we will see in Proposition 4.3.

**Remark 4.2.** Equation (4.3) in Notation 4.1 corresponds to (1.8).

**Proposition 4.3.** (1) If  $f \in \mathcal{D}_L$ , then  $f^2 \in \mathcal{D}_L$  and  $Lf^2 = \sigma^2 f'^2 + 2fLf$ .

(2)

$$Lh = 0, \quad Lh^2 = \sigma^2 h'^2.$$

**Proof.** (1) We observe  $f^2 \in \mathcal{D}_L$  because  $(f^2)' = 2ff' = (2f\phi) \exp(-\Sigma)$ . From Notation 4.1 (1) and the fact that  $\phi_2 := 2f\phi \in C^1$ , we conclude  $f^2 \in \mathcal{D}_L$ . By (4.3),

$$\begin{aligned} Lf^2 &= \phi_2' \exp(-\Sigma) \frac{\sigma^2}{2} = (f\phi)' \exp(-\Sigma) \sigma^2 \\ &= f' \sigma^2 \exp(-\Sigma) \phi + f \phi' \exp(-\Sigma) \sigma^2 = f'^2 \sigma^2 + 2fLf. \end{aligned}$$

(2) The proof follows by setting  $\phi = 1$ , using item (1) above and Notation 4.1 (1).  $\square$

We now formulate a standing assumption.

**Assumption 4.4.** •  $\Sigma$  given by (4.1) is a well-defined function.

• We assume the nonexplosion condition

$$\int_{-\infty}^0 e^{-\Sigma(x)} dx = \int_0^{+\infty} e^{-\Sigma(x)} dx = \infty. \quad (4.5)$$

**Remark 4.5.** (1) Under Assumption 4.4, the  $L$ -harmonic function  $h : \mathbb{R} \rightarrow \mathbb{R}$  defined in (4.4) is a  $C^1$ -diffeomorphism. In particular,  $h$  is surjective.



- (2) It is easy to verify that Assumption 4.4 implies the nonexplosion condition (3.16) in Proposition 3.13 in [12]. In [12], one could allow  $h$  not to be surjective.

**Remark 4.6.** When  $\sigma$  and  $b'$  are continuous functions, then  $\mathcal{D}_L = C^2$ . Indeed, in this case,  $\Sigma \in C^1$  and then  $f' = \exp(-\Sigma)\phi \in C^1$ . In particular,  $Lf$  corresponds to the classical definition.

**Remark 4.7.** We assume that Assumption 4.4 is satisfied. Let  $g \in \mathcal{D}_L$  be a fixed diffeomorphism of class  $C^1$  such that  $g' > 0$ . We set

$$\sigma_0^g := (\sigma g') \circ g^{-1}, \quad b^g = ((Lg) \circ g^{-1}) \quad (4.6)$$

and consider

$$L^g v := \frac{1}{2}(\sigma_0^g)^2 v'' + b^g v', \quad v \in \mathcal{D}_{L^g},$$

where we define  $\mathcal{D}_{L^g}$  according to Notation 4.1 replacing  $L$  with  $L^g$ .

By Remark 4.6, since  $L^g$  has continuous coefficients,  $\mathcal{D}_{L^g} = C^2$ .

**Proposition 4.8.** *We assume that Assumption 4.4 is satisfied. Let  $g \in \mathcal{D}_L$  be a fixed diffeomorphism of class  $C^1$  such that  $g' > 0$ . Then,  $f \in \mathcal{D}_L$  if and only if  $f \circ g^{-1}$  belongs to  $\mathcal{D}_{L^g}$  and*

$$Lf \circ g^{-1} = L^g(f \circ g^{-1}).$$

**Proof of Proposition 4.8.** By Notation 4.1, there exists  $\phi^g \in C^1$  such that

$$g' = \exp(-\Sigma)\phi^g. \quad (4.7)$$

Concerning the direct implication, if  $f \in \mathcal{D}_L$ , first we prove that  $f \circ g^{-1} \in \mathcal{D}_{L^g}$ . Again by Notation 4.1 there exists  $\phi^f \in C^1$  such that  $f' = \exp(-\Sigma)\phi^f$ . So,  $(f \circ g^{-1})' = \frac{f'}{g'} \circ g^{-1} = \frac{\phi^f}{\phi^g} \circ g^{-1} \in C^1$  because  $g^{-1} \in C^1$  and  $\phi^g > 0$ ; then,  $f \circ g^{-1} \in C^2$ . Note that, by Remark 4.7,  $\mathcal{D}_{L^g} = C^2$  and hence  $f \circ g^{-1} \in \mathcal{D}_{L^g}$ . Moreover, according to Notation 4.1 (2),

$$(\phi^f)' = \frac{2Lf}{\sigma^2} \exp(\Sigma), \quad (\phi^g)' = \frac{2Lg}{\sigma^2} \exp(\Sigma).$$

A direct calculation gives

$$(f \circ g^{-1})'' = \left( \frac{\phi^f}{\phi^g} \circ g^{-1} \right)' = \left[ \frac{2Lf}{g'^2 \sigma^2} - \frac{2Lg f'}{\sigma^2 g'^3} \right] \circ g^{-1}.$$

Consequently,

$$\frac{(\sigma_0^g)^2}{2} (f \circ g^{-1})'' = \left[ Lf - \frac{Lg f'}{g'} \right] \circ g^{-1}. \quad (4.8)$$

By (4.8),  $Lf \circ g^{-1} = \frac{(\sigma_0^g)^2}{2} (f \circ g^{-1})'' + (Lg) \circ g^{-1} (f \circ g^{-1})' = L^g(f \circ g^{-1})$ .

Let us discuss the converse implication. Suppose that  $f \circ g^{-1}$  belongs to  $\mathcal{D}_{L^g} = C^2$ . Again, according to Notation 4.1 (1), we need to show that  $f' \exp(\Sigma) \in$

$C^1$  which is equivalent to showing that  $(f' \exp(\Sigma)) \circ g^{-1}$  belongs to  $C^1$ . If  $\phi^g \in C^1$  is such that  $g' = \exp(-\Sigma)\phi^g$  (see (4.7)) we have

$$(f' \exp(\Sigma)) \circ g^{-1} = \left( f' \frac{\phi^g}{g'} \right) \circ g^{-1} = (f \circ g^{-1})'(\phi^g \circ g^{-1}),$$

which obviously belongs to  $C^1$ . Therefore  $f \in \mathcal{D}_L$ .  $\square$

By setting  $h = g$  in Proposition 4.8 we obtain the following corollary.

**Corollary 4.9.** *Let  $h$  be the function defined by (4.4). Then,  $f \in \mathcal{D}_L$  if and only if  $f \circ h^{-1} \in C^2$ . Moreover, by setting  $\varphi = f \circ h^{-1}$  for  $f \in \mathcal{D}_L$ , we have*

$$L(\varphi \circ h) \circ h^{-1} = L^h(\varphi) = \frac{1}{2}\sigma_0^2\varphi'',$$

where

$$\sigma_0 := \sigma_0^h = (\sigma h') \circ h^{-1}. \quad (4.9)$$

In [12], the authors also show that the existence and uniqueness of the solution to the martingale problem are conditioned to a nonexplosion feature. The proposition below is an easy consequence of Proposition 3.13 in [12], which concerns the well-posedness of the martingale problem with respect to  $L$  in the case  $\Gamma = 0$ .

**Proposition 4.10.** *Let  $v = \delta_{x_0}$ ,  $x_0 \in \mathbb{R}$  and suppose that Assumption 4.4 holds true. Then, the existence and uniqueness hold for the martingale problem related to  $L$  (i.e. with  $\Gamma = 0$ ) with respect to  $\mathcal{D}_L$  with initial condition  $v$ .*

**Remark 4.11.** By Proposition 3.2 of [12], if  $\Gamma = 0$  and  $(X, \mathbb{P})$  is a solution of the martingale problem given in Proposition 4.10, then there exists a  $\mathbb{P}$ -Brownian motion  $W$  such that (3.4) equals

$$\int_0^t (f'\sigma)(X_s) dW_s, \quad t \in [0, T].$$

## 4.2 The path-dependent framework

Let  $\sigma$  and  $b$  be functions in  $C(\mathbb{R})$  with  $\sigma > 0$  and  $\Gamma$  as defined in (1.2). Let us assume again Assumption 4.4 and let  $h$  be the function defined in (4.4). We recall that  $\sigma_0$  was defined in (4.9).

The first result explains how to reduce our path-dependent martingale problem to a path-dependent SDE.

**Proposition 4.12.** *Let  $X$  be a stochastic process on a probability space  $(\Omega, \mathcal{F}, \mathbb{P})$ .*

- (1)  *$(X, \mathbb{P})$  solves the martingale problem related to (3.3) with respect to  $\mathcal{D}_L$  if and only if the process  $Y = h(X)$  is a solution (with respect to  $\mathbb{P}$ ) of*

$$Y_t = Y_0 + \int_0^t \sigma_0(Y_s) dW_s + \int_0^t h'(h^{-1}(Y_s))\Gamma(s, h^{-1}(Y^s)) ds \quad (4.10)$$

for some  $\mathbb{P}$ -Brownian motion  $W$ .

(2) Let  $W$  be a Brownian motion on  $(\Omega, \mathcal{F}, \mathbb{P})$ . Then,  $X$  is a solution to the strong martingale problem with respect to  $\mathcal{D}_L$  and  $W$  if and only if (4.10) holds.

**Proof.** (1) We start proving the direct implication. According to (3.4) and the notations introduced therein,

$$M_t^h = h(X_t) - h(X_0) - \int_0^t Lh(X_s)ds - \int_0^t h'(X_s)\Gamma(s, X^s)ds \quad (4.11)$$

is a  $\mathbb{P}$ -local martingale on some measurable space  $(\Omega, \mathcal{F})$ .

In particular, by Proposition 4.3,  $Y$  satisfies

$$Y_t = Y_0 + \int_0^t h'(h^{-1}(Y_s))\Gamma(s, h^{-1}(Y^s))ds + M_t^h,$$

where  $M^h$  is a local martingale and hence  $Y$  is a semimartingale. We need now to evaluate  $[M^h]_t = [Y]_t$ . We apply (3.4) to  $f = h^2$  and, again by Proposition 4.3, we get

$$Y_t^2 = Y_0^2 + \int_0^t \sigma_0^2(Y_s)ds + 2 \int_0^t Y_s h'(h^{-1}(Y_s))\Gamma(s, h^{-1}(Y^s))ds + M_t^{h^2}, \quad (4.12)$$

where  $M^{h^2}$  is a local martingale, and we recall that  $\sigma_0$  was defined in (4.6). By integration by parts,

$$\begin{aligned} [Y]_t &= Y_t^2 - Y_0^2 - 2 \int_0^t Y_s dY_s \\ &= Y_t^2 - Y_0^2 + M_t - 2 \int_0^t Y_s h'(h^{-1}(Y_s))\Gamma(s, h^{-1}(Y^s))ds, \end{aligned}$$

where  $M_t = -2 \int_0^t Y_s dM_s^h$ . Therefore

$$Y_t^2 = Y_0^2 - M_t + 2 \int_0^t Y_s h'(h^{-1}(Y_s))\Gamma(s, h^{-1}(Y^s))ds + [Y]_t. \quad (4.13)$$

The semimartingale  $Y^2$  admits the two decompositions (4.12) and (4.13). By uniqueness,  $-M = M^{h^2}$  and  $\int_0^t \sigma_0^2(Y_s)ds = [Y]_t$ . By (4.11),

$$[M^h]_t = [Y]_t = \int_0^t \sigma_0^2(Y_s)ds.$$

Setting

$$W_t := \int_0^t \frac{dM_s^h}{\sigma_0(Y_s)}, \quad t \geq 0,$$

we have

$$[W]_t \equiv t.$$

Therefore, by Lévy's characterization of Brownian motion,  $W$  is a standard Brownian motion. Since

$$M^h = \int_0^\cdot \sigma_0(Y_s) dW_s,$$

(4.11) shows that  $Y$  solves (4.10).

Next, we prove the converse implication. Suppose that  $Y = h(X)$  satisfies (4.10) for some  $\mathbb{P}$ -Brownian motion  $W$ . We take  $f \in \mathcal{D}_L$ . By Corollary 4.9 we have  $\varphi \equiv f \circ h^{-1} \in C^2$ . Using Itô's formula and again Corollary 4.9, we get

$$\begin{aligned} \varphi(Y_t) &= \varphi(Y_0) + \int_0^t \varphi'(Y_s) dY_s + \frac{1}{2} \int_0^t \varphi''(Y_s) d[Y]_s \\ &= \varphi(Y_0) + \int_0^t \varphi'(Y_s) \sigma_0(Y_s) dW_s + \frac{1}{2} \int_0^t \varphi''(Y_s) \sigma_0^2(Y_s) ds \\ &= \varphi(Y_0) + \int_0^t \varphi'(Y_s) \sigma_0(Y_s) dW_s + \frac{1}{2} \int_0^t L^h \varphi(Y_s) \sigma_0^2(Y_s) ds \\ &= f(X_0) + \int_0^t Lf(X_s) ds + \int_0^t f'(X_s) \sigma(X_s) dW_s \\ &\quad + \int_0^t f'(X_s) \Gamma(s, X^s) ds. \end{aligned}$$

Therefore

$$\begin{aligned} f(X_t) - f(X_0) - \int_0^t Lf(X_s) ds - \int_0^t f'(X_s) \Gamma(s, X^s) ds \\ = \int_0^t f'(X_s) \sigma(X_s) dW_s \end{aligned}$$

is a local martingale, which concludes the proof.

(2) The converse implication follows in the same way as for item (1). The proof of the direct implication follows directly by Itô's formula.  $\square$

**Corollary 4.13.** *Let  $(X, \mathbb{P})$  be a solution of the martingale problem related to (3.3) with respect to  $\mathcal{D}_L$ . Then  $X$  is a Dirichlet process (with respect to its canonical filtration) and  $[X]_t = \int_0^t \sigma^2(X_s) ds$ ,  $t \in [0, T]$ .*

**Proof.** By Remark 4.5,  $h$  is a diffeomorphism. By Proposition 4.12,  $X = h^{-1}(Y)$ , where  $Y$  is obviously a semimartingale such that  $[Y]_t = \int_0^t \sigma_0^2(Y_s) ds$ ,  $t \in [0, T]$ . Consequently, by Remark 2.1,  $X$  is indeed a Dirichlet process and

$$\begin{aligned} [X]_t &= \int_0^t ((h^{-1})')^2 \sigma_0^2(Y_s) ds = \int_0^t \frac{\sigma_0^2}{(h' \circ h^{-1})^2}(Y_s) ds \\ &= \int_0^t \sigma^2(X_s) ds, \quad t \in [0, T]. \end{aligned}$$

$\square$

**Remark 4.14.** If  $X$  is a solution of the strong martingale problem with respect to  $\mathcal{D}_L$  and some Brownian motion  $W$ , then  $X$  is a Dirichlet process with respect to the canonical filtration of the Brownian motion.

An immediate consequence of Proposition 4.12 is the following corollary.

**Corollary 4.15.** *Suppose that  $\Gamma = 0$  and let  $(X, \mathbb{P})$  be a solution to the martingale problem related to  $L$  with respect to  $\mathcal{D}_L$ . Then,  $Y = h(X)$  is an  $\mathfrak{F}$ -local martingale where  $\mathfrak{F}$  is the canonical filtration of  $X$  with quadratic variation  $[Y] = \int_0^\cdot \sigma_0^2(Y_s) ds$ .*

### 4.3 Existence

We fix here the same conventions as in Section 4.2. In the sequel, we introduce the map  $\tilde{\Gamma} : \Lambda \rightarrow \mathbb{R}$  defined by

$$\tilde{\Gamma}(s, \eta) = \frac{\Gamma(s, \eta)}{\sigma(\eta(s))}, \quad (s, \eta) \in \Lambda. \quad (4.14)$$

At this point, we introduce the following technical assumption, which is in particular verified if  $\Gamma$  is bounded and  $\sigma = 1$ .

**Assumption 4.16.** *There is  $K > 0$  such that*

$$\sup_{s \in [0, T]} |\tilde{\Gamma}(s, h^{-1} \circ \eta^s)| \leq K \left( 1 + \sup_{s \in [0, T]} |\eta(s)| \right), \quad \forall \eta \in C([0, T]).$$

**Remark 4.17.** Let  $X$  be a stochastic process and set  $Y = h(X)$ . If Assumption 4.16 is fulfilled, then

$$\sup_{s \in [0, T]} |\tilde{\Gamma}(s, X^s)| \leq K \left( 1 + \sup_{s \in [0, T]} |Y_s| \right). \quad (4.15)$$

In particular  $\int_0^T \tilde{\Gamma}^2(s, X^s) ds < \infty$  a.s.

Proposition 4.18 below is a well-known extension of Novikov's criterion. It is an easy consequence of Corollary 5.14 in [16, Chapter 3].

**Proposition 4.18.** *Suppose Assumption 4.16 holds true. Let  $W$  be a Brownian motion and let  $X$  be a continuous and adapted process for which there exists a subdivision  $0 = t_0 < t_1 < \dots < t_n = T$  such that*

$$\mathbb{E} \left[ \exp \left( \frac{1}{2} \int_{t_{i-1}}^{t_i} |\tilde{\Gamma}(s, X^s)|^2 ds \right) \right] < \infty$$

for every  $i \in \{1, \dots, n\}$ . Then, the process

$$N_t = \exp \left( \int_0^t \tilde{\Gamma}(s, X^s) dW_s - \frac{1}{2} \int_0^t |\tilde{\Gamma}(s, X^s)|^2 ds \right)$$

is a martingale.

Next, we need a slight adaptation of the Dambis–Dubins–Schwarz theorem to the case of a finite interval. For the sake of completeness, we give the details here.

**Proposition 4.19.** *Let  $M$  be a local martingale vanishing at zero such that  $[M]_t = \int_0^t A_s ds$ ,  $t \in [0, T]$ . Then, on a possibly enlarged probability space, there exists a copy of  $M$  (still denoted by the same letter  $M$ ) with the same law and a Brownian motion  $\beta$  such that*

$$M_t = \beta \int_0^t A_s ds, \quad t \in [0, T].$$

**Proof.** Let us define

$$\tilde{M}_t = \begin{cases} M_t, & t \in [0, T], \\ M_T + B_t - B_T, & t > T, \end{cases}$$

where  $B$  is a Brownian motion independent of  $M$ . If the initial probability space is not rich enough, one considers an enlarged probability space containing a copy of  $M$  (still denoted by the same letter) with the same law and the independent Brownian motion  $B$ . Note that  $\tilde{M}$  is a local martingale with quadratic variation given by

$$[\tilde{M}]_t = \begin{cases} [M]_t, & t \in [0, T], \\ t - T + [M]_T, & t > T. \end{cases}$$

Observe that  $\lim_{t \rightarrow \infty} [\tilde{M}]_t = \infty$ . By the classical Dambis–Dubins–Schwarz theorem there exists a standard Brownian motion  $\beta$  such that a.s.  $\tilde{M}_t = \beta \int_0^t A_s ds$ ,  $t \geq 0$ . In particular,

$$M_t = \beta \int_0^t A_s ds, \quad 0 \leq t \leq T.$$

□

The proposition below is an adaptation of a well-known argument for Markov diffusions.

**Proposition 4.20.** *Suppose that Assumption 4.16 holds and that  $\sigma_0$  is bounded. Let  $(X, \mathbb{P})$  be a solution to the martingale problem related to (3.3) with respect to  $\mathcal{D}_L$  with  $\Gamma = 0$ . Let  $M^X$  be the local martingale component of  $X$ . We set*

$$W_t := \int_0^t \frac{1}{\sigma(X_s)} dM^X, \quad t \in [0, T].$$

Then

$$\exp\left(\int_0^t \tilde{\Gamma}(s, X^s) dW_s - \frac{1}{2} \int_0^t |\tilde{\Gamma}(s, X^s)|^2 ds\right), \quad t \in [0, T], \quad (4.16)$$

is a martingale.

**Remark 4.21.** Let  $(X, \mathbb{P})$  be a solution to the martingale problem related to (3.3) with respect to  $\mathcal{D}_L$  with  $\Gamma = 0$ . We recall that, by Corollary 4.13,  $X$  is an  $\mathfrak{F}$ -Dirichlet process ( $\mathfrak{F}$  being the canonical filtration) and  $[X] = [M^X] = \int_0^\cdot \sigma^2(X_s) ds$  so that, by Lévy’s characterization theorem,  $W$  is an  $\mathfrak{F}$ -Brownian motion.

**Proof of Proposition 4.20.** Let  $Y = h(X)$ . By Proposition 4.12, we know that  $[Y] = \int_0^\cdot \sigma_0^2(Y_s) ds$ . Let us choose  $k \geq |\sigma_0|_\infty^2 T$  and a subdivision  $\{t_0 = 0, \dots, t_n = T\}$  of  $[0, T]$  in such way that

$$c_i := \frac{3}{2}(t_i - t_{i-1})K^2 k < \frac{1}{2} \quad (4.17)$$

for  $i \in \{1, \dots, n\}$ . Here,  $K$  comes from Assumption 4.16. By (4.15), we know that

$$\int_{t_{i-1}}^{t_i} |\tilde{\Gamma}(s, X^s)|^2 ds \leq (t_i - t_{i-1})K^2 \left(1 + \sup_{s \in [0, T]} |Y_s|\right)^2 \quad (4.18)$$

for every  $i \in \{1, \dots, n\}$ . We set  $M_t = Y_t - Y_0$ ,  $t \in [0, T]$ , and we note that

$$\left(1 + \sup_{s \in [0, T]} |Y_s|\right)^2 \leq 3 \sup_{s \in [0, T]} |M_s|^2 + 3(1 + Y_0^2). \quad (4.19)$$

We recall that  $Y_0$  is deterministic. By applying Proposition 4.18, taking into account (4.18) and (4.19), we get

$$\begin{aligned} & \mathbb{E} \left( \exp \left( \frac{1}{2} \int_{t_{i-1}}^{t_i} \tilde{\Gamma}^2(s, X^s) ds \right) \right) \\ & \leq \mathbb{E} \left( \exp \left( \frac{3(t_i - t_{i-1})K^2}{2} \sup_{s \in [0, T]} |M_s|^2 \right) \right) \exp \left( \frac{3}{2} (t_i - t_{i-1})K^2 (1 + Y_0^2) \right). \end{aligned} \quad (4.20)$$

Since  $M$  is a local martingale vanishing at zero, Proposition 4.19 states that there is a copy (with the same distribution) of  $M$  (still denoted by the same letter) on another probability space, and a Brownian motion  $\beta$ , such that previous expression gives

$$\begin{aligned} & \mathbb{E} \left( \exp \left( \frac{3(t_i - t_{i-1})K^2}{2} \sup_{s \in [0, T]} |\beta_{[M]_s}|^2 \right) \right) \exp \left( \frac{3}{2} (t_i - t_{i-1})K^2 (1 + Y_0^2) \right) \\ & \leq \mathbb{E} \left( \exp \left( \frac{3(t_i - t_{i-1})K^2}{2} \sup_{\tau \in [0, k]} |\beta_\tau|^2 \right) \right) \exp \left( \frac{3}{2} (t_i - t_{i-1})K^2 (1 + Y_0^2) \right), \end{aligned} \quad (4.21)$$

the latter inequality being valid because  $[M]_t = \int_0^t \sigma_0^2(Y_s) ds$ . By (4.17) we get

$$\begin{aligned} \mathbb{E} \left[ \exp \left( \frac{1}{2} \int_{t_{i-1}}^{t_i} \tilde{\Gamma}^2(s, X^s) ds \right) \right] & \leq \mathbb{E} \left[ \exp \left( \frac{c_i}{k} \sup_{\tau \in [0, k]} |B_\tau|^2 \right) \right] \exp \left( \frac{c_i}{k} (1 + Y_0^2) \right) \\ & \leq \mathbb{E} \left[ \sup_{\tau \in [0, k]} \exp \left( \frac{c_i}{k} |B_\tau|^2 \right) \right] \exp \left( \frac{c_i}{k} (1 + Y_0^2) \right) \end{aligned} \quad (4.22)$$

for every  $i \in \{1, \dots, n\}$ . By Remark 4.22 below, for each  $i \in \{1, \dots, n\}$ , we have

$$\mathbb{E} \left[ \exp \left( \frac{c_i}{k} |B_\tau|^2 \right) \right] \leq \mathbb{E} [\exp(c_i G^2)] < \infty,$$

where  $G$  is a standard Gaussian random variable. Since  $x \mapsto \exp(\frac{c_i}{2k}x)$  is increasing and convex, and  $(|B_\tau|^2)_{\tau \geq 0}$  is a nonnegative square integrable submartingale,  $(\exp(\frac{c_i}{2k}|B_\tau|^2))$  is also a nonnegative submartingale. Consequently, by Doob's inequality (with  $p = 2$ ) the expectation on the right-hand side of (4.22) is finite. Finally by Proposition 4.18, (4.16) is a martingale.  $\square$

**Remark 4.22.** Let  $G$  be a standard Gaussian random variable. If  $c < \frac{1}{2}$  then

$$\mathbb{E}[\exp(cG^2)] < \infty.$$

Proposition 4.20 opens the way to the following existence result for the path-dependent martingale problem.

**Theorem 4.23.** *Suppose that Assumption 4.4 holds and that one of the two conditions below are fulfilled.*

- (1)  $\tilde{\Gamma}$  is bounded.
- (2)  $\tilde{\Gamma}$  fulfills Assumption 4.16 and  $\sigma_0$  is bounded.

Then existence holds for the martingale problem related to (3.3) with respect to  $\mathcal{D}_L$ .

**Proof.** By Proposition 4.10, there is a solution  $(X, \mathbb{P})$  to the above-mentioned martingale problem with  $\Gamma = 0$ . By Remark 4.11, there is a Brownian motion  $W$  such that

$$f(X_t) - f(X_0) - \int_0^t Lf(X_s)ds = \int_0^t (f'\sigma)(X_s)dW_s \quad (4.23)$$

for every  $f \in \mathcal{D}_L$ . We define the process

$$V_t := \exp\left(\int_0^t \tilde{\Gamma}(s, X^s)dW_s - \frac{1}{2} \int_0^t \tilde{\Gamma}^2(s, X^s)ds\right).$$

Under item (1),  $V$  is a martingale by Novikov's condition. Under item (2), Proposition 4.20 says that  $V$  is a martingale. We define

$$\tilde{W}_t := W_t - \int_0^t \tilde{\Gamma}(s, X^s)ds. \quad (4.24)$$

By Girsanov's theorem, (4.24) is a Brownian motion under the probability  $\mathbb{Q}$  such that

$$d\mathbb{Q} := \exp\left(\int_0^T \tilde{\Gamma}(s, X^s)dW_s - \frac{1}{2} \int_0^T \tilde{\Gamma}^2(s, X^s)ds\right)d\mathbb{P}.$$

Applying (4.24) to (4.23), we obtain

$$f(X_t) - f(X_0) - \int_0^t Lf(X_s)ds - \int_0^t f'(X_s)\Gamma(s, X^s)ds = \int_0^t (f'\sigma)(X_s)d\tilde{W}_s$$

for every  $f \in \mathcal{D}_L$ . Since  $\int_0^t (f'\sigma)(X_s)d\tilde{W}_s$  is a local martingale under  $\mathbb{Q}$ ,  $(X, \mathbb{Q})$  is proved to be a solution to the martingale problem in the statement.  $\square$

#### 4.4 Uniqueness

We use here again the notation  $\tilde{\Gamma}$  introduced in (4.14).

**Proposition 4.24.** *Suppose that Assumption 4.4 is satisfied. Then uniqueness holds for the martingale problem related to (3.3) with respect to  $\mathcal{D}_L$ .*



**Proof.** Let  $(X^i, \mathbb{P}^i)$ ,  $i = 1, 2$ , be two solutions of the martingale problem related to (3.3) with respect to  $\mathcal{D}_L$ . Let us fix  $i = 1, 2$ . By Corollary 4.13,  $X^i$  is an  $\mathfrak{F}^{X^i}$ -Dirichlet process with respect to  $\mathbb{P}^i$ , such that  $[X^i] \equiv \int_0^\cdot \sigma(X_s^i)^2 ds$ . Let  $M^i$  be the martingale component of  $X^i$ . Since  $[M^i] \equiv \int_0^\cdot \sigma(X_s^i)^2 ds$ , by Lévy's characterization theorem, the process

$$W_t^i = \int_0^t \frac{dM_s^i}{\sigma(X_s^i)}, \quad t \in [0, T], \quad (4.25)$$

is an  $\mathfrak{F}^{X^i}$ -Brownian motion. In particular,  $W^i$  is a Borel functional of  $X^i$ .

By means of localization (similarly to Proposition 5.3.10 in [16]), without loss of generality we can suppose  $\tilde{\Gamma}$  to be bounded. We define the process (whose random variables are also Borel functionals of  $X^i$ )

$$V_t^i = \exp\left(-\int_0^t \tilde{\Gamma}(s, X^{i,s}) dW_s^i - \frac{1}{2} \int_0^t (\tilde{\Gamma}(s, X^{i,s}))^2 ds\right),$$

which, by Novikov's condition, is a  $\mathbb{P}^i$ -martingale. This allows us to define the probability  $\mathbb{Q}^i$  such that  $d\mathbb{Q}^i = V_T^i d\mathbb{P}^i$ .

By Girsanov's theorem, under  $\mathbb{Q}^i$ ,  $B_t^i := W_t^i + \int_0^t \tilde{\Gamma}(s, X^{i,s}) ds$  is a Brownian motion. Therefore,  $(X^i, \mathbb{Q}^i)$  solves the martingale problem related to  $L$  ( $\Gamma = 0$ ) with respect to  $\mathcal{D}_L$ . By uniqueness of the martingale problem with respect to  $\mathcal{D}_L$  and  $\Gamma = 0$  (see Proposition 4.10),  $X^i$  (under  $\mathbb{Q}^i$ ),  $i = 1, 2$ , have the same law. Hence, for every Borel set  $B \in \mathfrak{B}(C[0, T])$ , we have

$$\mathbb{P}^1\{X^1 \in B\} = \int_\Omega \frac{1}{V_T^1(X^1)} 1_{\{X^1 \in B\}} d\mathbb{Q}^1 = \int_\Omega \frac{1}{V_T^2(X^2)} 1_{\{X^2 \in B\}} d\mathbb{Q}^2 = \mathbb{P}^2\{X^2 \in B\}.$$

Therefore,  $X^1$  under  $\mathbb{P}^1$  has the same law as  $X^2$  under  $\mathbb{P}^2$ . Finally, uniqueness holds for the martingale problem related to (3.3) with respect to  $\mathcal{D}_L$ .  $\square$

#### 4.5 Results on pathwise uniqueness

Before exploring conditions for strong existence and uniqueness for the martingale problem, we state and prove Proposition 4.27, which constitutes a crucial preliminary step.

Let  $\bar{\Gamma} : \Lambda \rightarrow \mathbb{R}$  be a generic Borel functional. Related to it, we formulate the following technical assumption.

**Assumption 4.25.** (1) *There exists a function  $l : \mathbb{R}_+ \rightarrow \mathbb{R}_+$  such that  $\int_0^\epsilon l^{-2}(u) du = \infty$  for all  $\epsilon > 0$  and*

$$|\sigma_0(x) - \sigma_0(y)| \leq l(|x - y|).$$

(2)  *$\sigma_0$  has at most linear growth.*

(3) *There exists  $K > 0$  such that*

$$|\bar{\Gamma}(s, \eta^1) - \bar{\Gamma}(s, \eta^2)| \leq K \left( |\eta^1(s) - \eta^2(s)| + \int_0^s |\eta^1(r) - \eta^2(r)| dr \right)$$

*for all  $(s, \eta^1), (s, \eta^2) \in \Lambda$ .*

$$(4) \bar{\Gamma}_\infty := \sup_{s \in [0, T]} |\bar{\Gamma}(s, 0)| < \infty.$$

**Remark 4.26.** One typical example of nonanticipative functional which satisfies (3) is given by

$$\bar{\Gamma}(s, \eta) = \Phi\left(s, \eta(s), \int_0^s \eta(r) dr\right),$$

where  $\Phi : [0, T] \times \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$  is a Lipschitz real function in the second and third variables.

**Proposition 4.27.** *Suppose that Assumption 4.25 is satisfied and fix  $y_0 \in \mathbb{R}$ . Then, pathwise uniqueness holds for the SDE with dynamics*

$$Y_t = y_0 + \int_0^t \sigma_0(Y_s) dW_s + \int_0^t \bar{\Gamma}(s, Y^s) ds, \quad (4.26)$$

i.e.  $E(\sigma_0, 0, \bar{\Gamma})$ .

The proof of the Proposition 4.27 generalizes the techniques of the Yamada–Watanabe pathwise uniqueness theorem for Markovian SDEs (see, e.g., Theorem 5.2.19 in [16]). Before proceeding with that proof, we state a lemma which is an easy consequence of Problem 5.3.15 in [16].

**Lemma 4.28.** *Suppose the assumptions in Proposition 4.27 are in force. Let  $Y$  be a solution of (4.26) and let  $m \geq 2$  be an integer. Then, there exists a constant  $C > 0$ , depending on the linear growth constant of  $\sigma_0$ ,  $Y_0$ ,  $T$ ,  $m$ , and the quantities  $(K, \bar{\Gamma}_\infty)$  given in Assumptions 4.25 (3)–(4), such that*

$$\mathbb{E}\left(\sup_{t \leq T} |Y_t|^m\right) \leq C.$$

**Proof of Proposition 4.27.** Let  $Y^1, Y^2$  be two solutions on the same probability space with respect to the same Brownian motion  $W$  of (4.26) such that  $Y_0^1 = Y_0^2 = y_0$ . In the sequel, we set  $\Delta_t = Y_t^1 - Y_t^2$ ,  $t \in [0, T]$ . By Lemma 4.28, we have

$$\mathbb{E}\left(\sup_{t \in [0, T]} |Y_t^i|^2\right) < \infty \quad (4.27)$$

for  $i = 1, 2$ . By the assumption on  $\sigma_0$ , this obviously gives

$$\mathbb{E}\left(\int_0^T |\sigma_0(Y_t^i)|^2 dt\right) < \infty \quad (4.28)$$

for  $i = 1, 2$ . We observe

$$\Delta_t = \int_0^t (\bar{\Gamma}(s, Y^{1,s}) - \bar{\Gamma}(s, Y^{2,s})) ds + \int_0^t (\sigma_0(Y_s^1) - \sigma_0(Y_s^2)) dW_s, \quad t \in [0, T]. \quad (4.29)$$

We recall from the proof of Proposition 2.13 in [16, Chapter 5], the existence of the functions

$$\Psi_n(x) = \int_0^{|x|} \int_0^y \rho_n(u) du dy,$$

such that for every  $x \in \mathbb{R}$

$$0 \leq \rho_n(x) \leq \frac{2}{nl^2(x)}, \quad |\Psi'_n(x)| \leq 1, \quad |\Psi_n(x)| \leq |x|, \quad \lim_{n \rightarrow \infty} \Psi_n(x) = |x|. \quad (4.30)$$

By applying Itô's formula and using (4.29), we get

$$\begin{aligned} \Psi_n(\Delta_t) &= \int_0^t \Psi'_n(\Delta_s) [\bar{\Gamma}(s, Y^{1,s}) - \bar{\Gamma}(s, Y^{2,s})] ds \\ &\quad + \frac{1}{2} \int_0^t \Psi''_n(\Delta_s) [\sigma_0(Y_s^1) - \sigma_0(Y_s^2)]^2 ds \\ &\quad + \int_0^t \Psi'_n(\Delta_s) [\sigma_0(Y_s^1) - \sigma_0(Y_s^2)] dW_s. \end{aligned}$$

By using Assumption 4.25 and (4.30), we get

$$\Psi_n(\Delta_t) \leq \int_0^t K \left( |Y_s^1 - Y_s^2| + \int_0^s |Y_r^1 - Y_r^2| dr \right) ds + \frac{t}{n} + M_t, \quad (4.31)$$

where  $M_t = \int_0^t \Psi'_n(\Delta_s) [\sigma_0(Y_s^1) - \sigma_0(Y_s^2)] dW_s$  is a local martingale. Since  $\Psi'_n$  is bounded, the estimate (4.28) ensures that  $M$  is a (even square integrable) martingale.

We take the expectation, and applying Fubini's theorem in (4.31), obtain

$$\mathbb{E} \Psi_n(\Delta_t) \leq K \int_0^t \mathbb{E} |Y_s^1 - Y_s^2| ds + KT \int_0^t \mathbb{E} |Y_s^1 - Y_s^2| ds + \frac{t}{n}, \quad (4.32)$$

since  $\mathbb{E} M_t = 0$ . Passing to the limit when  $n \rightarrow \infty$ , by Lebesgue's dominated convergence theorem, we get

$$\mathbb{E} |\Delta_t| \leq (K + TK) \int_0^t \mathbb{E} |\Delta_s| ds. \quad (4.33)$$

By Gronwall's inequality, we obtain  $\mathbb{E} |\Delta_t| = 0$ . By the continuity of the sample paths of  $Y^1, Y^2$  we conclude that  $Y^1, Y^2$  are indistinguishable.  $\square$

We come back to the framework of the beginning of Section 4.1. We assume again the validity of Assumption 4.4. We recall the definition of the harmonic function  $h$  defined by  $h(0) = 0$ ,  $h'(x) = e^{-\Sigma}$ , see (4.4). We recall the notation  $\sigma_0 = (\sigma h') \circ h^{-1}$ . We define

$$\bar{\Gamma}(s, \eta) := h'(h^{-1}(\eta(s))) \Gamma(s, h^{-1}(\eta^s)), \quad s \in [0, T], \quad \eta \in C([0, T]). \quad (4.34)$$

**Corollary 4.29.** *Suppose that Assumptions 4.4 and 4.25 (related to  $\bar{\Gamma}$  introduced in (4.34)) are fulfilled. Then uniqueness holds for the martingale problem related to (3.3) with respect to  $\mathcal{D}_L$ .*

**Proof.** It follows by using Assumptions 4.4 and 4.25 on  $\bar{\Gamma}$  and applying Proposition 4.27 and Proposition 4.12.  $\square$

**Theorem 4.30.** *Suppose that Assumptions 4.4 and 4.25 (related to  $\bar{\Gamma}$  introduced in (4.34)) and one of two hypotheses below are in force:*

- (1)  $\tilde{\Gamma}$  defined in (4.14) is bounded,
- (2)  $\sigma_0$  is bounded and uniformly elliptic.

*Then strong existence and pathwise uniqueness hold for the SDE with dynamics (4.10).*

**Proof.** By Proposition 4.27, pathwise uniqueness holds. Indeed, taking into account the expression of (4.34), equation (4.10) is a particular case of (4.26).

In order to prove existence, we will apply Theorem 4.23. For this purpose, we need to verify that either Hypothesis (1) or (2) of that theorem hold. Hypothesis (1) in the above statement coincides with Hypothesis (1) in Theorem 4.23. Assume the validity of (2) in the above statement, and we check that Assumption 4.16 holds true. By (4.14) and the definition of  $\bar{\Gamma}$  in (4.34), we obtain

$$\sigma_0(\eta(s))\tilde{\Gamma}(s, (h^{-1} \circ \eta)) = \bar{\Gamma}(s, \eta). \quad (4.35)$$

The fact that  $\frac{1}{\sigma_0}$  is bounded jointly with (3) in Assumption 4.25 yields the existence of a constant  $K_1 > 0$  such that

$$|\tilde{\Gamma}(s, (h^{-1} \circ \eta))| \leq K_1 \left( |\eta(s)| + \int_0^s |\eta(r)| dr \right) + |\bar{\Gamma}(s, 0)|.$$

By (4) in Assumption 4.25 and the simple estimate

$$\int_0^s |\gamma(r)| dr \leq s \sup_{r \in [0, s]} |\gamma(r)|,$$

we can conclude that Assumption 4.16 holds.

By Theorem 4.23, existence holds for the martingale problem related to (3.3) with respect to  $\mathcal{D}_L$ , and by Proposition 4.12 (1), we have that (4.10) has a solution. At this point, we can apply the Yamada–Watanabe theorem to guarantee that the solution is actually strong. We remark that the Yamada–Watanabe theorem (in the path-dependent case) proof is the same as the one in the Markovian case, which is for instance stated in Proposition 3.20 [16, Chapter 5].  $\square$

As a consequence of Proposition 4.12 and Theorem 4.30, we obtain the following result.

**Corollary 4.31.** *Under the same assumptions as in Theorem 4.30, strong existence and pathwise uniqueness hold for the martingale problem related to (3.3) with respect to  $\mathcal{D}_L$ .*

## A Different notions of solutions when $b'$ is a function

Let us suppose below that  $\sigma, b' : \mathbb{R} \rightarrow \mathbb{R}$  are locally bounded Borel functions and  $\Gamma$  is as given in (1.2). As already mentioned, for simplicity we will only consider deterministic initial conditions  $x_0$ .

**Definition A.1.** Let  $(\Omega, \mathcal{F}, \mathbb{P})$  be a probability space,  $(W_t)_{t \geq 0}$  be a Brownian motion and  $x_0 \in \mathbb{R}$ . A solution  $X$  of  $E(\sigma, b', \Gamma; x_0)$  (depending on the probability space  $(\Omega, \mathcal{F}, \mathbb{P})$  and the Brownian motion  $W$ ) is an  $\mathfrak{F}^W$ -progressively measurable process fulfilling (3.1), with  $\xi = x_0$  a.s.

**Definition A.2** (Strong existence). We say that **strong existence** holds for equation  $E(\sigma, b', \Gamma)$  if for any probability space  $(\Omega, \mathcal{F}, \mathbb{P})$  carrying a Brownian motion  $(W_t)_{t \geq 0}$  and  $x_0 \in \mathbb{R}$ , there exists a process  $(X_t)_{t \geq 0}$  which is a solution to  $E(\sigma, b', \Gamma; x_0)$ .

**Definition A.3** (Pathwise uniqueness). We say that pathwise uniqueness holds for equation  $E(\sigma, b', \Gamma)$  if the following property is fulfilled for every  $x_0 \in \mathbb{R}$ . Let  $(\Omega, \mathcal{F}, \mathbb{P})$  be a probability space carrying a Brownian motion  $(W_t)_{t \geq 0}$ . If two processes  $X, \tilde{X}$  are two solutions to  $E(\sigma, b', \Gamma; x_0)$ , then  $X$  and  $\tilde{X}$  are indistinguishable.

**Definition A.4** (Existence in law). Let  $x_0 \in \mathbb{R}$ . We say that existence in law holds for  $E(\sigma, b', \Gamma; x_0)$  if there exists a probability space  $(\Omega, \mathcal{F}, \mathbb{P})$  carrying a Brownian motion  $(W_t)_{t \geq 0}$  and a process  $(X_t)_{t \geq 0}$  such that  $(X, \mathbb{P})$  is a solution of  $E(\sigma, b', \Gamma; x_0)$ , see Definition A.1.

We say that existence in law holds for  $E(\sigma, b', \Gamma)$  if existence in law holds for  $E(\sigma, b', \Gamma; x_0)$ , for every  $x_0 \in \mathbb{R}$ .

**Definition A.5** (Uniqueness in law). Let  $x_0 \in \mathbb{R}$ . We say that **uniqueness in law** holds for  $E(\sigma, b', \Gamma; x_0)$  if we have the following. Suppose we have a probability space  $(\Omega, \mathcal{F}, \mathbb{P})$  (respectively  $(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{\mathbb{P}})$ ) carrying a Brownian motion  $(W_t)_{t \geq 0}$  (respectively  $(\tilde{W}_t)_{t \geq 0}$ ). We suppose that a process  $(X_t)_{t \geq 0}$  (resp. a process  $(\tilde{X}_t)_{t \geq 0}$ ) is a solution of  $E(\sigma, b', \Gamma; x_0)$ , such that both  $X_0 = x_0, \mathbb{P}$  a.s. and  $\tilde{X}_0 = x_0, \tilde{\mathbb{P}}$  a.s. Uniqueness in law means that  $X$  and  $\tilde{X}$  must have the same law as random elements taking values in  $C([0, T])$  or  $C(\mathbb{R}_+)$ .

We say that **uniqueness in law holds** for  $E(\sigma, b', \Gamma)$  if uniqueness in law holds for  $E(\sigma, b', \Gamma; x_0)$  for every  $x_0 \in \mathbb{R}$ .

## Acknowledgement

The authors are very grateful to both referees for their very careful reading of the first submitted version and for their relevant comments which allowed to drastically improve the paper.

## Funding

The research related to this paper was financially supported by the Regional Program MATH-AmSud 2018 grant 88887.197425/2018-00. A.O. acknowledges the financial support of CNPq Bolsa de Produtividade de Pesquisa grant 303443/2018-9.

## References

- [1] Bass, R.F., Chen, Z.-Q.: Stochastic differential equations for Dirichlet processes. *Probab. Theory Relat. Fields* **121**(3), 422–446 (2001). [MR1867429. https://doi.org/10.1007/s004400100151](https://doi.org/10.1007/s004400100151)

- [2] Bass, R.F., Chen, Z.-Q.: Brownian motion with singular drift. *Ann. Probab.* **31**(2), 791–817 (2003). [MR1964949](#). <https://doi.org/10.1214/aop/1048516536>
- [3] Blei, S., Engelbert, H.J.: One-dimensional stochastic differential equations with generalized and singular drift. *Stoch. Process. Appl.* **123**(12), 4337–4372 (2013). [MR3096356](#). <https://doi.org/10.1016/j.spa.2013.06.014>
- [4] Brox, Th.: A one-dimensional diffusion process in a Wiener medium. *Ann. Probab.* **14**(4), 1206–1218 (1986). [MR0866343](#)
- [5] Cannizzaro, G., Chouk, K.: Multidimensional SDEs with singular drift and universal construction of the polymer measure with white noise potential. *Ann. Probab.* **46**(3), 1710–1763 (2018). 05. [MR3785598](#). <https://doi.org/10.1214/17-AOP1213>
- [6] Castrequini, R.A., Russo, F.: Path dependent equations driven by Hölder processes. *Stoch. Anal. Appl.* **37**(3), 480–498 (2019). [MR3946622](#). <https://doi.org/10.1080/07362994.2019.1585263>
- [7] Delarue, F., Diel, R.: Rough paths and 1d SDE with a time dependent distributional drift: application to polymers. *Probab. Theory Relat. Fields* **165**, 1–63 (2016). [MR3500267](#). <https://doi.org/10.1007/s00440-015-0626-8>
- [8] Dereudre, D., Roelly, S.: Path-dependent infinite-dimensional SDE with non-regular drift: An existence result. *Ann. Inst. Henri Poincaré Probab. Stat.* **53**(2), 641–657 (2017). [MR3634268](#). <https://doi.org/10.1214/15-AIHP728>
- [9] Engelbert, H.-J., Schmidt, W.: On one-dimensional stochastic differential equations with generalized drift. In: *Stochastic differential systems, Marseille-Luminy, 1984. Lecture Notes in Control and Inform. Sci.*, vol. 69, pp. 143–155. Springer, Berlin (1985). [MR0798317](#). <https://doi.org/10.1007/BFb0005069>
- [10] Engelbert, H.-J., Wolf, J.: Strong Markov local Dirichlet processes and stochastic differential equations. *Teor. Veroâtn. Primen.* **43**(2), 331–348 (1998). [MR1679006](#). <https://doi.org/10.4213/tvp1468>
- [11] Flandoli, F., Issoglio, E., Russo, F.: Multidimensional SDEs with distributional coefficients. *Trans. Am. Math. Soc.* **369**, 1665–1688 (2017). [MR3581216](#). <https://doi.org/10.1090/tran/6729>
- [12] Flandoli, F., Russo, F., Wolf, J.: Some SDEs with distributional drift. I. General calculus. *Osaka J. Math.* **40**(2), 493–542 (2003). [MR1988703](#)
- [13] Flandoli, F., Russo, F., Wolf, J.: Some SDEs with distributional drift. II. Lyons-Zheng structure, Itô’s formula and semimartingale characterization. *Random Oper. Stoch. Equ.* **12**(2), 145–184 (2004). [MR2065168](#). <https://doi.org/10.1163/156939704323074700>
- [14] Frikha, N., Li, L.: Weak uniqueness and density estimates for SDEs with coefficients depending on some path-functionals. *ArXiv preprint* (2019). [1707.01295](#). [MR4076774](#). <https://doi.org/10.1214/19-AIHP992>
- [15] Hu, Y., Shi, Z., Yor, M.: Rates of convergences of diffusions with drifted Brownian potentials. *Transl. Am. Math. Soc.* **351**(10), 3915–3934 (1999). [MR1637078](#). <https://doi.org/10.1090/S0002-9947-99-02421-6>
- [16] Karatzas, I., Shreve, S.E.: *Brownian motion and stochastic calculus*, 2nd edn. Graduate Texts in Mathematics, vol. 113. Springer, New York (1991). [MR1121940](#). <https://doi.org/10.1007/978-1-4612-0949-2>
- [17] Mohammed, S-E.A.: *Stochastic functional differential equations*, vol. 99. Pitman Advanced Publishing Program (1984). [MR0754561](#)
- [18] Ohashi, A., Russo, F., Teixeira, A.: SDEs for Bessel processes in low dimension and path-dependent extensions (2020). Preprint.

- [19] Portenko, N.I.: Generalized diffusion processes. Translations of Mathematical Monographs, vol. 83. American Mathematical Society, Providence, RI (1990). Translated from the Russian by H. H. McFaden. [MR1104660](#). <https://doi.org/10.1090/mmono/083>
- [20] Rogers, L.C.G., Williams, D.: Diffusions, Markov processes, and Martingales. Cambridge University Press (2000). [MR1780932](#). <https://doi.org/10.1017/CBO9781107590120>
- [21] Russo, F., Trutnau, G.: About a construction and some analysis of time inhomogeneous diffusions on monotonely moving domains. *J. Funct. Anal.* **221**(1), 37–82 (2005). [MR2124897](#). <https://doi.org/10.1016/j.jfa.2004.08.007>
- [22] Russo, F., Trutnau, G.: Some parabolic PDEs whose drift is an irregular random noise in space. *Ann. Probab.* **35**(6), 2213–2262 (2007). [MR2353387](#). <https://doi.org/10.1214/009117906000001178>
- [23] Russo, F., Vallois, P.: The generalized covariation process and Itô formula. *Stoch. Process. Appl.* **59**(1), 81–104 (1995). [MR1350257](#). [https://doi.org/10.1016/0304-4149\(95\)93237-A](https://doi.org/10.1016/0304-4149(95)93237-A)
- [24] Scheutzow, M.: Stochastic delay equations. Lecture Notes, CIMPA School (2018)
- [25] Seignourel, P.: Discrete schemes for processes in random media. *Probab. Theory Relat. Fields* **118**(3), 293–322 (2000). [MR1800534](#). <https://doi.org/10.1007/PL00008743>
- [26] Stroock, D.W., Varadhan, S.R.S.: Multidimensional diffusion processes. *Classics in Mathematics*. Springer, Berlin (2006). Reprint of the 1997 edition. [MR2190038](#)
- [27] Zvonkin, A.K.: A transformation of the phase space of a diffusion process that will remove the drift. *Mat. Sb.* **93**(135), 129–149 (1974). 152. [MR0336813](#)