

Multidimensional compound Poisson approximations for symmetric distributions

Vydas Čekanavičius^{a,*}, Simona Jokubauskienė^b

^a*Institute of Applied Mathematics, Faculty of Mathematics and Informatics, Vilnius University, Naugarduko 24, Vilnius 03225, Lithuania*

^b*Department of Mathematics and Statistics, Faculty of Informatics, Vytautas Magnus University, Universiteto 10-202, 53361, Akademija, Lithuania*

vydas.ceknavicius@mif.vu.lt (V. Čekanavičius), simona.jokubauskiene@vdu.lt (S. Jokubauskienė)

Received: 29 January 2026, Revised: 13 May 2026, Accepted: 14 May 2026,
Published online: 26 May 2026

Abstract Distribution of the sum of independent identically distributed symmetric lattice vectors is approximated by the accompanying compound Poisson law and the second-order Hipp-type signed compound Poisson measure. Bergström-type asymptotic expansion is constructed. The accuracy of approximation is estimated in the total variation metric and, in many cases, is of the order $O(n^{-1})$.

Keywords Compound Poisson approximation, convolution of distributions, sums of random vectors, total variation metric, the first uniform Kolmogorov theorem

2020 MSC 60F99, 60G50

1 Introduction

Compound Poisson (CP) approximations have many applications and are one of the popular fields of research in modern probability theory, see, for example, [2, 3, 8, 9, 15, 23, 24] and the references therein. Research papers in this field are numerous. However, the absolute majority of results are proved for one-dimensional random variables. The situation is very different for random vectors (r.v.s), where results are a few and usually obtained only for Kolmogorov, Prokhorov, or local metrics, see

*Corresponding author.

[10, 12, 27] and the references therein. Note that, for sequences of r.v.s, a significant part of multidimensional research is devoted to normal approximation [4, 6] or behavior of tails of distributions [16]. We are primarily interested in triangular array of vectors.

In this paper, for sums of lattice identically distributed symmetric r.v.s, we adapt Le Cam's convolution approach and properties of metrics to prove results for total variation. Apart from the general case, when no moment assumptions are imposed, we consider mixtures of distributions concentrated on separate lines with finite analogues of variances, demonstrating how known one-dimensional results can be applied to obtain the multidimensional estimates.

We introduce necessary notation. The sets of all integers and natural numbers are denoted respectively by \mathbb{Z} and \mathbb{N} . By \mathcal{M}_d and $\mathcal{F}_d \subset \mathcal{M}_d$ we denote the set of all finite (signed) d -dimensional measures and the set of all d -dimensional distributions, respectively. Further on $d < \infty$ is some fixed natural number. Let $I_{\vec{a}} \in \mathcal{F}_d$ be the distribution concentrated at $\vec{a} \in \mathbb{R}^d$, $I = I_{\vec{0}}$, where $\vec{0} = (0, 0, \dots, 0) \in \mathbb{R}^d$. Let $M, V \in \mathcal{M}_d$, then their convolution, for any d -dimensional Borel set A is defined as $(M * V) \{A\} = \int_{\mathbb{R}^d} M \{A - \vec{x}\} V \{d\vec{x}\}$.

By $\mathcal{L}(\vec{X})$ we denote the distribution of a r.v. \vec{X} . Note that if $S_n = \vec{X}_1 + \vec{X}_2 + \dots + \vec{X}_n$, where all r.v.s. are independent identically distributed (i.i.d.) and $\mathcal{L}(\vec{X}_1) = F$, then $\mathcal{L}(S_n) = F^{*n}$, where power is understood in the convolution sense. In this paper we prefer measure notation over r.v.s, since the main tools used for the proofs are properties of norms and convolutions.

Exponential measure for $M \in \mathcal{M}_d$ is defined as $\exp(M) = \sum_{j \geq 0} M^{*j} / j!$, where $M^{*0} = I$.

Compound Poisson distribution with parameter $\lambda > 0$ and compounding distribution $F \in \mathcal{F}_d$ is defined as $\exp(\lambda(F - I))$. Note that CP distribution can be expressed also as a distribution of random sum $\vec{Y} = \sum_{j=0}^{\pi_\lambda} \vec{X}_j$, where $\vec{X}, \vec{X}_1, \vec{X}_2, \dots$ are i.i.d. r.v.s independent from Poisson random variable π_λ and $\mathcal{L}(\vec{X}) = F$. CP distribution $\exp(F - I)$ is called *accompanying distribution* to $F \in \mathcal{F}_d$. If $F \in \mathcal{F}_d$, but $\lambda < 0$, then $\exp(\lambda(F - I))$ is called signed compound Poisson measure (SCP). All CP distributions are infinitely divisible.

For $F, G \in \mathcal{F}_d$ total variation distance and total variation norm are defined as

$$d_{TV}(F, G) := \sup_B |F \{B\} - G \{B\}| =: \frac{1}{2} \|F - G\|_{TV},$$

where supremum is taken over all d -dimensional Borel sets. Total variation norm can also be defined via Hahn decomposition, see [8], p. 228. For $F, G \in \mathcal{F}_d$ with $\text{supp } F, \text{supp } G \subset \mathbb{Z}^d$

$$\|F - G\|_{TV} = \sum_{\vec{m} \in \mathbb{Z}^d} |F \{\vec{m}\} - G \{\vec{m}\}|.$$

To make proofs more concise the same symbol C is used to denotes all positive constants. If ambiguities can arise we supply C with indices. Symbol $C(N)$ denotes quantity, which depends only on N . Notation $f(n) = O(n^{-k})$ means that $f(n)n^k$ is bounded by some C for all n . Multiplication is superior to division. For any $x \in \mathbb{R}$ let $[x]$ denote the integer part of x .

2 Known results

General CP approximations for independent symmetric vectors in Kolmogorov and Prokhorov metrics were investigated in [10, 28] (see also references, therein). In this paper, we consider stronger total variation metric. Multidimensional estimates in total variation for CP approximations are few. Dependent vectors were investigated in Theorem 6.8 [17]. Non-symmetric vectors were considered by Roos [20–22], who used the so-called Kerstan's method. In [14], Kerstan's method was applied for symmetric vectors concentrated on coordinate axes of \mathbb{Z}^d . Set $\bar{e}_1 = (1, 0, \dots, 0), \dots, \bar{e}_m = (0, \dots, 0, 1, 0, \dots, 0), \bar{e}_d = (0, \dots, 0, 1)$. Let $\vec{X}_1, \dots, \vec{X}_n$ be independent r.v.s taking values in \mathbb{Z}^d and let for $m = 1, 2, \dots, d, i = 1, 2, \dots, n$:

$$\begin{aligned} \mathcal{L}(\vec{X}_i) &= q_i I + \sum_{m=1}^d p_{i,m} F_m, \quad q_i = \mathbb{P}(\vec{X}_i = \vec{0}), \quad q_i + \sum_{m=1}^d p_{i,m} = 1, \\ \text{supp } F_m &\subset \{k\bar{e}_m, k = \pm 1, \pm 2, \dots\}, \quad F_m \{k\bar{e}_m\} = F_m \{-k\bar{e}_m\}, \\ \lambda_m &= \sum_{i=1}^n p_{i,m}, \quad \sigma_m^2 = 2 \sum_{k=1}^{\infty} k^2 F_m \{k\bar{e}_m\}, \quad 0 \leq q_i, p_{i,m} \leq 1, \\ \alpha &= \sum_{i=1}^n g(2(1 - q_i)) \min \left\{ 2^{-3/2} \sum_{m=1}^d \frac{p_{i,m}^2}{\lambda_m}, \left(\sum_{m=1}^d p_{i,m} \right)^2 \right\}, \\ g(x) &= 2 \sum_{s=2}^{\infty} \frac{x^{s-2}}{s!} (s-1) = \frac{2e^x(e^{-x} - 1 + x)}{x^2}, \quad x \neq 0. \end{aligned}$$

Observe that in above setting coordinates of \vec{X}_i are uncorrelated and, for example, $\mathbb{P}(\vec{X}_i = (1, 1, 1, \dots, 1)) = 0$. In [14] it was proved that if $2\alpha\epsilon < 1, \sigma_m^2 < \infty$ ($m = 1, 2, \dots, d$), then

$$\begin{aligned} d_{TV} \left(\prod_{i=1}^{*n} \mathcal{L}(\vec{X}_i), \prod_{i=1}^{*n} \exp(\mathcal{L}(\vec{X}_i) - I) \right) \\ \leq \frac{15.98}{(1 - 2\alpha\epsilon)^{3/2}} \sum_{l=1}^d (1 + \sigma_l) \sum_{m=1}^d \frac{1}{\lambda_m^2} \sum_{i=1}^n p_{i,m}^2. \end{aligned} \quad (1)$$

If $d, \sigma_m, p_{i,m} \asymp C$, then the rate of accuracy in (1) is $O(n^{-1})$.

In the case of i.i.d r.v.s. $\vec{X}_i \equiv \vec{X}$, when $p_{i,m} \equiv p_m$ and $q = \mathbb{P}(\vec{X} = \vec{0}) \geq 4/5$, it was proved that

$$\begin{aligned} d_{TV} \left(\left(qI + \sum_{m=1}^d p_m F_m \right)^{*n}, \exp \left\{ n \sum_{m=1}^d p_m (F_m - I) \right\} \right) \\ \leq \frac{17.34}{n} \left(\sum_{m=1}^d \sqrt{1 + \sigma_m} \right)^2, \end{aligned} \quad (2)$$

see [14], Theorem 3.2. Kerstan's method has obvious restrictions. For (1) and (2) to hold, probabilities $\mathbb{P}(\vec{X}_i = \vec{0})$ must be large. Consider, for example, the case $d = 1$,

$q_i = p_i = 1/2$. Then $2\alpha e > 1$ and (1) cannot be applied. Moreover, σ_m can be large even if $\text{supp } \vec{X}_i$ is finite. The assumption that all coordinate vectors are uncorrelated is also very restrictive.

Note that any distribution with $q = F\{\vec{0}\} > 0$ can be written as $F = qI + (1-q)V$. From one-dimensional Poisson approximation to the binomial law it follows then that

$$d_{TV}(F^{*n}, \exp(n(F-I))) \leq \min(1-q, n(1-q)^2), \quad (3)$$

see [8], p. 207 and (3.15) for details. However, (3) does not take into account the symmetry of F and requires q to be very close to 1, much closer than needed for (2).

3 Results

3.1 The general case of vectors with possibly correlated coordinates

We will demonstrate that for identically distributed symmetric r.v.s, the assumptions used for (2) can be significantly weakened. Let

$$\vec{X} \sim F \in \mathcal{F}_d^s, \quad \text{supp } F \subset \mathbb{Z}^d, \quad q := F\{\vec{0}\} = \mathbb{P}(\vec{X} = \vec{0}) \in (0, 1). \quad (4)$$

Assumptions (4) are more general than used in (1), (2). Indeed, in (2) it was required that $q \geq 4/5$, meanwhile in (4) q can be reasonably small. Moreover, non-zero probabilities are not restricted to coordinate axes. Further let $p_{\vec{x}} := \mathbb{P}(\vec{X} = \vec{x}) = F\{\vec{x}\}$ and let $\sum_{\vec{x}}$ denote the summation over all $\vec{x} \in \text{supp } F \setminus \{\vec{0}\}$. Let

$$\delta(y) = \sum_{\vec{x}} \min(1, 4yp_{\vec{x}}). \quad (5)$$

Note that $\delta(y) \leq 4y(1-q)$, $\delta(y/k) \leq \delta(y)$ if $k \geq 1$ and $\delta(y) \leq 2N$ if $\text{supp } F = \{\vec{0}, \pm \vec{x}_1, \dots, \pm \vec{x}_N\}$.

The following theorem gives a general idea about what constant can be expected for large n .

Theorem 1. *Let assumption (4) hold. Then, for any $n \in \mathbb{N}$*

$$d_{TV}(F^{*n}, \exp(n(F-I))) \leq \frac{\delta^2(n/2)}{4n} \left(1 + \frac{C(\delta(nq) + 1)}{q^4 \sqrt{n}}\right) + \frac{C(1-q)^{9/2}}{q^5 n \sqrt{n}}. \quad (6)$$

The following shorter version of Theorem 1 with the smaller power of q in denominator holds.

Theorem 1*. *Let assumption (4) hold. Then, for any $n \in \mathbb{N}$*

$$d_{TV}(F^{*n}, \exp(n(F-I))) \leq \frac{C(1 + \delta^2(n))}{nq^{7/2}}. \quad (7)$$

Theorem 1 can be used even if (1) and (2) are not applicable.

Example 1. Let $d = 2$ and $\vec{X}, \vec{X}_1, \dots, \vec{X}_n$ be i.i.d. random vectors with

$$\begin{aligned} \mathbb{P}(\vec{X} = (0, 0)) &= \frac{1}{2}, & \mathbb{P}(\vec{X} = (\pm k, 0)) &= \frac{1}{2k(k+1)(k+2)}, & k = 1, 2, \dots \\ \mathbb{P}(\vec{X} = (0, \pm j)) &= \frac{1}{2j(j+1)(j+2)}, & j &= 1, 2, \dots \end{aligned}$$

Observe that (2) is not applicable, since $q = 1/2 < 4/5$ and $\sigma_1, \sigma_2 = \infty$. On the other hand,

$$\delta(n) \leq 2 \left(2 \sum_{k=1}^{\lfloor n^{1/3} \rfloor + 1} 1 + ne \sum_{k=\lfloor n^{1/3} \rfloor + 2}^{\infty} \frac{1}{k(k+1)(k+2)} \right) < 10n^{1/3}$$

and from (7) follows estimate $d_{TV}(F^{*n}, \exp(n(F-I))) \leq Cn^{-1/3}$.

Next we use SCP approximation of Hipp's type, see [11, 21] and [24], Ch. 10.7. Let for any real a

$$D := \exp \left((F-I) - \frac{(F-I)^{*2}}{2} \right), \quad D^{*a} := \exp \left(a(F-I) - \frac{a(F-I)^{*2}}{2} \right) \quad (8)$$

Theorem 2. Let assumption (4) hold. Then, for any $n \in \mathbb{N}$,

$$d_{TV}(F^{*n}, D^{*n}) \leq \frac{15.75\delta^3(nq/6)}{q^{7/2}n^2} \left(1 + \frac{C}{q^{7/2}\sqrt{n}} \right) + \frac{C(1-q)^{15/2}}{q^8 n^2 \sqrt{n}}. \quad (9)$$

SCP approximation D can be viewed as first order asymptotic expansion in the exponent. Next we consider more usual first order asymptotic expansion to accompanying CP distribution.

Theorem 3. Let assumption (4) hold. Then, for any $n \in \mathbb{N}$

$$\begin{aligned} d_{TV} \left(F^{*n}, \exp(n(F-I)) * \left(I - \frac{n(F-I)^{*2}}{2} \right) \right) \\ \leq \frac{C(\delta^3(nq) + \delta^4(nq))}{q^{11/2}n^2} + \frac{C(1-q)^6}{q^{13/2}n^2}. \end{aligned} \quad (10)$$

It is easy to check that for the Example 1 the accuracy in (9) (respectively (10)) is of the order $O(n^{-1})$ (respectively $O(n^{-2/3})$).

It is known that in Kolmogorov metric the closeness of subsequent convolutions F^{*n} and $F^{*(n+1)}$ can be estimated by $C(d)n^{-1}$, see [26]. Assumption (4) allows for similar estimate in total variation.

Theorem 4. Let assumption (4) hold. Then, for any $n \in \mathbb{N}$

$$d_{TV}(F^{*n}, F^{*(n+1)}) \leq \frac{C(\delta^2(n) + 1)}{q^{7/2}n}. \quad (11)$$

Orders of accuracy in (6)–(11) are explicit, when \vec{X} takes only finite number of values.

Corollary 1. *Let assumption (4) hold and let $\text{supp } F = \{\vec{0}, \pm \vec{x}_1, \pm \vec{x}_2, \dots, \pm \vec{x}_N\}$, $N \geq d$. Then*

$$\begin{aligned} d_{TV}(F^{*n}, \exp(n(F-I))) &\leq \frac{N^2}{n} \left(1 + \frac{CN}{q^5 \sqrt{n}}\right), \\ d_{TV}(F^{*n}, F^{*(n+1)}) &\leq CN^2 q^{-7/2} n^{-1}, \quad d_{TV}(F^{*n}, D^{*n}) \leq CN^3 q^{-7/2} n^{-2}, \\ d_{TV}(F^{*n}, \exp(n(F-I)) * (I - n(F-I)^{*2}/2)) &\leq CN^4 q^{-13/2} n^{-2}. \end{aligned}$$

Note that only the number of points matter, not their values. The following example shows that even for r.v.s with bounded supports estimate (6) can be more accurate than (2).

Example 2. Let $d = 2$, $\mathbb{P}(\vec{X} = (\pm 1, 0)) = \mathbb{P}(\vec{X} = (\pm n, 0)) = \mathbb{P}(\vec{X} = (0, \pm 1)) = \mathbb{P}(\vec{X} = (0, \pm n)) = 0.025$, $\mathbb{P}(\vec{X} = (0, 0)) = 0.8$. Then (2) gives trivial order of approximation $O(1)$. Meanwhile from Corollary 1 it follows that approximation by $\exp(n(F-I))$ is of the order $O(n^{-1})$.

Observe that estimates in Corollary 1 remain meaningful even if $q \rightarrow 0$, $N \rightarrow \infty$ slowly. For example, if $q = 1/\ln n$, $N = \ln n$ the estimates still have quite good order of approximation.

Though absolute constant in Corollary 1 is very reasonable, the following result shows that, at least in the scheme of sequences, it might be much smaller.

Proposition 1. *Let $\text{supp } F = \{\vec{0}, \pm \vec{x}_1, \pm \vec{x}_2, \dots, \pm \vec{x}_N\}$, $N \geq d$ and let $N, q, p_{\vec{x}}$ and support of F do not depend on n . Then*

$$\lim_{n \rightarrow \infty} n d_{TV}(F^{*n}, \exp(n(F-I))) \leq 0.17504N + 0.05855N(N-1). \quad (12)$$

Explicit constants can be obtained for the difference of two CP distributions with the same compounding distribution F .

Theorem 5. *Let assumption (4) hold and let $\text{supp } F = \{\vec{0}, \pm \vec{x}_1, \pm \vec{x}_2, \dots, \pm \vec{x}_N\}$, $N \geq d$. Then, for any $a, b > 0$, the following estimate holds*

$$d_{TV}(\exp(a(F-I)), \exp(b(F-I))) \leq \left(1 + \frac{2N+1}{e}\right) \frac{|b-a|}{\max(a, b)}.$$

If q is very large (F is almost degenerate) the following trivial estimate can be used

$$d_{TV}(\exp(a(F-I)), \exp(b(F-I))) \leq 2|b-a|(1-q).$$

So far we considered short asymptotic expansions. Next we demonstrate that, at least for D , it is possible to construct long so-called Bergström-type [5] asymptotic expansion. Let

$$M_k := F^{*n} - \sum_{m=0}^k \binom{n}{m} D^{*(n-m)} * (F-D)^{*m}, \quad k \leq n-1. \quad (13)$$

Theorem 6. *Let assumption (4) hold. Then, for any $n \in \mathbb{N}$, $k < n$, the following estimate holds*

$$\|M_k\|_{TV} \leq \frac{C(k)\delta^{3(k+1)}(nq)}{q^{6k+11/2}n^{2(k+1)}} + \frac{C(k)(1-q)^{6(k+1)}}{n^{2(k+1)}q^{6k+13/2}}.$$

Corollary 2. *Let assumption (4) hold and let $\text{supp } F = \{\vec{0}, \pm \vec{x}_1, \pm \vec{x}_2, \dots, \pm \vec{x}_N\}$, $N \geq d$. Then*

$$\|M_k\|_{TV} \leq \frac{C(k)N^{3(k+1)}}{q^{6k+13/2}n^{2(k+1)}}.$$

Theorem 6 can be used as intermediate result for construction of longer expansions.

3.2 The case of r.v.s concentrated on finite number of lines

Observe that for some cases estimate (2) is more accurate than the one in (6).

Example 3. Let $d = 2$ and $\vec{X}, \vec{X}_1, \dots, \vec{X}_n$ be i.i.d random vectors with

$$\mathbb{P}(\vec{X} = (0, 0)) = \frac{7}{9}, \quad \mathbb{P}(\vec{X} = (\pm k, 0)) = \frac{1}{2k(k+1)(k+2)(k+3)}, \quad k = 1, 2, \dots$$

$$\mathbb{P}(\vec{X} = (0, \pm j)) = \frac{1}{2j(j+1)(j+2)(j+3)}, \quad j = 1, 2, \dots$$

Then, arguing similarly to Example 1, we get

$$\delta(n) \leq 2 \left(2 \sum_{k=1}^{\lfloor n^{1/4} \rfloor + 1} 1 + 2ne \sum_{k=\lfloor n^{1/4} \rfloor + 2}^{\infty} \frac{1}{k(k+1)(k+2)(k+3)} \right) \leq Cn^{1/4}.$$

From (7) it follows that $d_{TV}(F^{*n}, \exp(n(F-I))) \leq Cn^{-1/2}$. Meanwhile, $\sigma_1^2, \sigma_2^2 \leq 2$ and from (2) it follows that $d_{TV}(F^{*n}, \exp(n(F-I))) \leq Cn^{-1}$.

Below we obtain some generalizations of (2). We assume that random vectors are concentrated on a finite number of lines and satisfy certain moment conditions. As before we assume that $\vec{X}, \vec{X}_1, \dots, \vec{X}_n$ are i.i.d. r.v.s in \mathbb{R}^d , $\mathcal{L}(\vec{X}) = F$. We also assume that

1. For some natural integer $K \geq d$

$$F = qI + \sum_{m=1}^K p_m F_m, \quad q, p_1, \dots, p_m \in (0, 1), \quad \sum_{m=1}^K p_m = 1 - q. \quad (14)$$

2. There exist non-zero vectors $\vec{y}_1, \vec{y}_2, \dots, \vec{y}_K \in \mathbb{R}^d$ such that

$$\text{supp } F_m \subset \{k\vec{y}_m : k \in \mathbb{Z} \setminus \{0\}\}, \quad \text{supp } F_m \cap \text{supp } F_k = \emptyset, \quad k \neq m. \quad (15)$$

3. All $\text{supp } F_m \neq \emptyset$ and F_m are symmetric:

$$F_m \{k\vec{y}_m\} = F_m \{-k\vec{y}_m\}, \quad k \in \{1, 2, \dots\}, \quad m = 1, \dots, K. \quad (16)$$

4. There exist k_m^0 such that

$$F_m\{k_m^0 \vec{y}_m\} > 0, \quad F_m\{(k_m^0 + 1)\vec{y}_m\} > 0. \quad (17)$$

Distribution considered in (2) is special case of above setting, since it suffices to take $K = d$ and $\vec{y}_m = \bar{e}_m$. We can define ‘projection’ of F to \vec{y}_m by considering one-dimensional integer-valued random variable Y such that

$$\mathbb{P}(Y = k) := F_m\{k\vec{y}_m\}, \quad k \in \mathbb{N}. \quad (18)$$

Then (17) is an assumption that Y is concentrated on a lattice with the maximal span equal to one. Though not formulated explicitly, assumption (17) was used for the proof of (2) in [14]. We denote the variance of Y by

$$\sigma_m^2 = \sum_{k=-\infty}^{\infty} k^2 F_m\{k\vec{y}_m\} = 2 \sum_{k=1}^{\infty} k^2 F_m\{k\vec{y}_m\}.$$

First we show that assumption $q \geq 4/5$ used in (2) can be dropped.

Theorem 7. *Let assumptions (14)–(17) hold. Then, for any $n \in \mathbb{N}$,*

$$\begin{aligned} & d_{TV}(F^{*n}, \exp(n(F-I))) \\ & \leq \frac{1.76}{n} \left(\sum_{m=1}^K \sqrt{1 + \sigma_m} \right)^2 \left(1 + \frac{C}{\sqrt{n} q^5} \sum_{m=1}^K \sqrt{1 + \sigma_m} \right). \end{aligned} \quad (19)$$

A shorter version with non-explicit constant can also be proved.

Theorem 7*. *Let assumptions (14)–(17) hold. Then, for any $n \in \mathbb{N}$,*

$$d_{TV}(F^{*n}, \exp(n(F-I))) \leq \frac{C}{nq^{7/2}} \left(\sum_{m=1}^K \sqrt{1 + \sigma_m} \right)^2. \quad (20)$$

Next we consider SCP approximation by D .

Theorem 8. *Let assumptions (14)–(17) hold. Then, for any $n \in \mathbb{N}$,*

$$d_{TV}(F^{*n}, D^{*n}) \leq \frac{C}{n^2 q^{13/2}} \left(\sum_{m=1}^K \sqrt{1 + \sigma_m} \right)^3. \quad (21)$$

We can formulate analogues of Theorems 3 and 4.

Theorem 9. *Let assumptions (14)–(17) hold. Then, for any $n \in \mathbb{N}$,*

$$d_{TV}(F^{*n}, \exp(n(F-I)) * (I - n(F-I)^{*2}/2)) \leq \frac{C}{n^2 q^{13/2}} \left(\sum_{m=1}^K \sqrt{1 + \sigma_m} \right)^4. \quad (22)$$

Theorem 10. *Let assumptions (14)–(17) hold. Then, for any $n \in \mathbb{N}$,*

$$d_{TV}(F^{*n}, F^{*(n+1)}) \leq \frac{C}{nq^{7/2}} \left(\sum_{m=1}^K \sqrt{1 + \sigma_m} \right)^2.$$

Table 1. $d_{TV}(F^{*n}, \exp(n(F - I)))$ for small n

p	$n=10$	$n=100$	$n=500$
0.01	0.0028088	0.0018381	0.0003544
0.10	0.0182749	0.0017669	0.0003509
0.25	0.0185827	0.0017586	0.0003503
0.45	0.0544728	0.0017514	0.0003503

Finally we formulate analogue of Theorem 5.

Theorem 11. *Let assumptions (14)–(17) hold. Then, for any $a, b > 0$,*

$$d_{TV}(\exp(a(F - I)), \exp(b(F - I))) \leq \frac{1.7|b - a|}{\max(a, b)} \sum_{m=1}^K \sqrt{1 + \sigma_m}.$$

Note that by assumption $K \geq d$. Therefore, strictly speaking, estimates in Theorems 7–11 depend on d and become trivial if $d \rightarrow \infty$.

3.3 Some numerical simulations

From the results of this paper it follows that the accuracy of approximation is very good for large n . We give some examples of estimates for simulated data when n is small. We begin from the case $d = 1$. Let $\mathbb{P}(X = -1) = \mathbb{P}(X = 1) = p$, $\mathbb{P}(X = 0) = 1 - 2p$, $F = \mathcal{L}(X)$, that is $\widehat{F}(t) = q + p(e^{-it} + e^{it})$.

Estimates for different simulations are given in Table 1. On one hand, we see that the accuracy of approximation is very good. On the other hand, if we consider $n \geq 100$, then the accuracy of approximation is $\approx 0.175/n$, which is better than $\approx 1/n$ from Corollary 1 and very close to the estimate in Proposition 1. Note also that, for $n = 10$ and small p , the accuracy of approximation is determined by (3).

Next let us consider direct extension of the previous simulation to 2-dimensional case. Let $q_i = 1 - 2p_i$, $\widehat{F}_i(t_i) = q_i + p_i(e^{it_i} + e^{-it_i})$, $i = 1, 2$, $\widehat{F}(t_1, t_2) = \widehat{F}_1(t_1)\widehat{F}_2(t_2)$. For approximation we use convolution of accompanying distributions $\widehat{H}(t_1, t_2) = \exp(\widehat{F}_1(t_1) - 1) \exp(\widehat{F}_2(t_2) - 1)$.

In Table 2 we present some simulated estimates for $d_{TV}(F^{*n}, H^{*n})$. From the triangle inequality $d_{TV}(F_1^{*n} * F_2^{*n}; \exp(n(F_1 - I)) * \exp(n(F_2 - I))) \leq d_{TV}(F_1^{*n}; \exp(n(F_1 - I))) + d_{TV}(F_2^{*n}; \exp(n(F_2 - I)))$ and the first simulation, one can expect the accuracy at least of the order $\approx 0.34/n$, $n \geq 100$. From Table 2 we see that, in reality, the accuracy is even better $\approx 0.25/n$.

Table 2. $d_{TV}(F^{*n}, H^{*n})$ for small n

p_1	p_2	$n=10$	$n=100$	$n=500$
0.01	0.01	0.0048421	0.0025247	0.0004952
0.10	0.10	0.0258073	0.0024872	0.0004974
0.10	0.45	0.0552711	0.0024914	0.0004976
0.45	0.45	0.0678198	0.0024953	0.0004978

4 Auxiliary results

First we formulate general facts about exponential measures. Let $M, V \in \mathcal{M}_d$, $k \in \mathbb{N}$. We have $\exp(M) * \exp(V) = \exp(M+V)$. From definition of exponential measure it follows that

$$e^M = I + M + \frac{M^{*2}}{2!} + \cdots + \frac{M^{*k}}{k!} + \frac{M^{*(k+1)}}{k!} * \int_0^1 e^{\tau M} (1-\tau)^k d\tau. \quad (23)$$

Let $F \in \mathcal{F}_d$, $a > 0$. The following well-known relations hold: $\|F\|_{TV} = 1$,

$$\|M * V\|_{TV} \leq \|M\|_{TV} \|V\|_{TV}, \quad \|\exp(M)\|_{TV} \leq \exp(\|M\|_{TV}), \quad (24)$$

$$\|(F - I)^{*k} * \exp(a(F - I))\|_{TV} \leq \|(F - I) * \exp(a(F - I)/k)\|_{TV}^k, \quad (25)$$

$$\left\| \sum_{j=0}^{\infty} \alpha_j F^{*j} \right\|_{TV} \leq \sum_{j=0}^{\infty} |\alpha_j| = \left\| \sum_{j=0}^{\infty} \alpha_j I_j \right\|_{TV}, \quad (26)$$

where $I_j \in \mathcal{F}_1$ is one-dimensional distribution concentrated at j . Sometimes inequality (26) is formulated in terms of random sums, see Eq. (3.1), [25]. It allows to reformulate many one-dimensional results for multidimensional case. Next lemma exemplifies such reformulation.

Lemma 1. *If $\lambda > 0$, $k \in \mathbb{N}$, $\vec{x} \in \mathbb{R}^d$. Then*

$$\|(I_{-\vec{x}} + I_{\vec{x}} - 2I) * \exp(\lambda(I_{-\vec{x}} + I_{\vec{x}} - 2I))\|_{TV} \leq \frac{1}{\lambda}, \quad (27)$$

$$\sup_{P \in \mathcal{F}_d} \|(P - I)^{*k} * e^{\lambda(P - I)}\|_{TV} \leq \left(\frac{2k}{e\lambda} \right)^{k/2}. \quad (28)$$

Estimate (27) follows from Lemma 4.6 in [7] and (26). Estimate (28) follows from Proposition A.2.7 in [1], (26) and (25).

The following technical estimates follow from Eq. (31) in [13] and from [19] Lemma 6.

Lemma 2. *Let $\lambda > 0$. Then*

$$\int_{-\pi}^{\pi} \sin^2(t/2) e^{-2\lambda \sin^2(t/2)} dt \leq \frac{\sqrt{\pi}}{\lambda^{3/2}}, \quad \int_{-\pi}^{\pi} e^{-2\lambda \sin^2(t/2)} dt \leq \frac{\pi}{\sqrt{3\lambda}} \left(1 + \sqrt{\frac{\pi}{2}} \right).$$

Next we present generalization of Lemma B.2 from [8].

Lemma 3. *Let $p_i \in (0, 1)$, ($i = 1, 2, \dots, n$), $p_0 = \max_i p_i$, $\tau \in [0, 1]$. Then*

$$\sup_{F \in \mathcal{F}_d} \left\| \exp \left(\frac{(1+p_0)}{2} \sum_{i=1}^n p_i (F - I) - \frac{\tau}{2} \sum_{i=1}^n p_i^2 (F - I)^{*2} \right) \right\|_{TV} \leq \frac{3.5}{\sqrt{1-p_0}}.$$

Particularly, if $p_i \equiv p$, then

$$\sup_{F \in \mathcal{F}_d} \left\| \exp \left(\frac{np(1+p)}{2} (F - I) - \frac{np^2\tau}{2} (F - I)^{*2} \right) \right\|_{TV} \leq \frac{3.5}{\sqrt{1-p}}. \quad (29)$$

Proof. In view of (26), it suffices to prove lemma for one-dimensional case $F = I_1 \in \mathcal{F}_1$. Let

$$M := \frac{(1+p_0)}{2} \sum_{i=1}^n p_i (I_1 - I) - \frac{\tau}{2} \sum_{i=1}^n p_i^2 (I_1 - I)^{*2}.$$

We denote by i complex unit, that is, $i^2 = -1$. Set $e^{ix} := \cos x + i \sin x$ and let $\widehat{M}(t)$ be Fourier transform of M . Set $a = \sum_{i=1}^n p_i$, $v = a(1+p_0)/2$ and

$$\widehat{M}_c(t) := \widehat{M}(t) - itv = \frac{a(1+p_0)}{2} (e^{it} - 1 - it) - \frac{\tau}{2} \sum_{i=1}^n p_i^2 (e^{it} - 1)^2,$$

$$b = 2.634 \sqrt{\frac{a}{1-p_0}}.$$

Then

$$\begin{aligned} \left| e^{\widehat{M}_c(t)} \right| &\leq \exp \left(-(1+p_0)a \sin^2(t/2) + \frac{p_0}{2} a |e^{it} - 1|^2 \right) \\ &\leq \exp \left(-(1-p_0)a \sin^2(t/2) \right). \end{aligned}$$

Observe that

$$\left| \widehat{M}'_c(t) \right| \leq \frac{a(1+p_0)}{2} |e^{it} - 1| + ap_0 |e^{it} - 1| \leq 4a |\sin(t/2)|.$$

Therefore,

$$\left| \left(e^{\widehat{M}_c(t)} \right)' \right|^2 \leq 16a^2 \sin^2(t/2) e^{-2(1-p_0)a \sin^2(t/2)}.$$

and by Lemma 2

$$b \int_{-\pi}^{\pi} \left| e^{\widehat{M}_c(t)} \right|^2 dt + \frac{1}{b} \int_{-\pi}^{\pi} \left| \left(e^{\widehat{M}_c(t)} \right)' \right|^2 dt \leq \frac{21.54}{1-p_0}.$$

If $b > 2.5$, then lemma's statement follows from formula of inversion (see (D.10) in [8])

$$\|e^M\|_{TV}^2 \leq \frac{1+b\pi}{2\pi} \int_{-\pi}^{\pi} \left(\left| e^{\widehat{M}_c(t)} \right|^2 + b^{-2} \left| \left(e^{\widehat{M}_c(t)} \right)' \right|^2 \right) dt$$

and a simple estimate

$$\frac{1+b\pi}{2\pi} \leq \frac{b}{2} \left(1 + \frac{2}{5\pi} \right) \leq 0.5637 b.$$

If $b \leq 2.5$, then $a \leq 0.901(1-p_0)$, $a(1+3p_0) \leq 0.901(1-p_0)(1+3p_0) \leq 1.2014$ and

$$\|e^M\|_{TV} \leq \exp \left(\frac{a(1+p_0)}{2} \|I_1 - I\|_{TV} + \frac{ap_0}{2} \|I_1 - I\|_{TV}^2 \right) \leq e^{a(1+3p_0)} \leq 3.33.$$

□

Lemma 4. Let $P \in \mathcal{F}_1$ be a symmetric distribution concentrated on $\mathbb{Z} \setminus \{0\}$. Assume that P has a finite variance σ^2 . Then for any $\lambda > 0$ and $j \in \mathbb{N}$

$$\|(P - I)^{*j} * e^{\lambda(P-I)}\|_{TV} \leq \frac{3.6\sqrt{1+\sigma} j^{j+1/4}}{\lambda^j e^j}$$

Lemma 4 is part of Lemma 4.6 from [7]. Similar bounds are valid for the compound binomial distribution. Next lemma is Lemma 4 in [18].

Lemma 5. Let $q = 1 - p$. If $0 < p < 1$, $j, n \in \mathbb{N}$, then

$$\sup_{F \in \mathcal{F}_d} \|(F - I)^{*j} * (qI + pF)^{*n}\|_{TV} \leq \sqrt{e} j^{1/4} \left(\frac{n}{n+j}\right)^{n/2} \left(\frac{j}{(n+j)pq}\right)^{j/2}. \quad (30)$$

We recall that notation Θ is used for all measures satisfying $\|\Theta\|_{TV} \leq 1$.

Lemma 6. Let $F = qI + pV \in \mathcal{F}_d$, $q = 1 - p \in (0, 1)$, $m \in \mathbb{N}$. Then for any $m \in \mathbb{N}$, $a \in \mathbb{R}$

$$D - I = C(F - I) * \Theta, \quad \|D^{*a}\|_{TV} \leq e^{4|a|}, \quad (31)$$

$$D^{*m} = \frac{3.5}{\sqrt{q}} \exp\left(\frac{mq(F - I)}{2}\right) * \Theta_m, \quad \|D^{*m}\|_{TV} \leq \frac{3.5}{\sqrt{q}}, \quad (32)$$

$$\begin{aligned} F - D &= \frac{(F - I)^{*3}}{3} + C_1(F - I)^{*4} * \Theta = C_2(F - I)^{*3} * \Theta \\ &= C_2 p^3 (V - I)^{*3} * \Theta. \end{aligned} \quad (33)$$

Proof. The first estimate in (32) follows from (29). The second estimate in (32) follows from the fact that the total variation norm of any CP distribution is equal to 1 and, therefore,

$$\|D^{*m}\|_{TV} \leq \frac{3.5}{\sqrt{q}} \|\exp(mq(F - I)/2)\|_{TV} \|\Theta_m\|_{TV} \leq \frac{3.5}{\sqrt{q}}.$$

Observe that $\|F - I\|_{TV} \leq \|F\|_{TV} + \|I\|_{TV} = 2$ and, therefore, by (24)

$$\|D^{*a}\|_{TV} \leq \exp(|a| \|F - I\|_{TV} + |a| \|F - I\|_{TV}^2 / 2) \leq e^{4|a|}.$$

Expressions (31) and (33) follow directly from definition of exponential measure. \square

The following result is of special interest, since it shows that long asymptotic expansions can be constructed without assumptions of symmetry. Let M_k be defined by (13).

Lemma 7. Let $F = qI + pV \in \mathcal{F}_d$, $q = 1 - p \in (0, 1)$, $k, n \in \mathbb{N}$, $k \leq n - 1$. Then

$$\|M_k\|_{TV} \leq \frac{C(k) p^{3(k+1)/2}}{n^{(k+1)/2} q^{(3k+4)/2}}.$$

Proof. If $n \leq 12$, then

$$\|M_k\|_{TV} \leq 1 + \sum_{m=0}^k \binom{n}{m} e^{4(n-m)} (1 + e^4)^m \leq 1 + (1 + 2e^4)^{12} = C = \frac{n^k C}{n^k} \leq \frac{12^k C}{n^k}.$$

Therefore, we further assume $n \geq 13$. Bergström identity [5] allows to write

$$M_k = \sum_{m=k+1}^n \binom{m-1}{k} F^{*(n-m)} * (F-D)^{*(k+1)} * D^{*(m-k-1)}.$$

Noting that

$$\sum_{m=k+1}^n \binom{m-1}{k} = \binom{n}{k+1} \quad (34)$$

and applying (33), (32), (28), (30), (34) we obtain

$$\begin{aligned} \|M_k\|_{TV} &\leq \sum_{m=k+1}^n \binom{m-1}{k} \|F^{*(n-m)} * (F-D)^{*(k+1)} * D^{*(m-1)}\|_{TV} \\ &\leq C(k) \sum_{m \leq \lfloor n/2 \rfloor} \binom{m-1}{k} \|(F-I)^{*3(k+1)} * F^{*(n-m)}\|_{TV} \|D^{*(m-k-1)}\|_{TV} \\ &\quad + C(k) \sum_{m > \lfloor n/2 \rfloor} \binom{m-1}{k} \|F^{*(n-m)}\|_{TV} \|(F-I)^{*3(k+1)} * D^{*(m-k-1)}\|_{TV} \\ &\leq \frac{C(k)n^{k+1}}{\sqrt{q}} p^{3(k+1)} \|(V-I)^{*3(k+1)} * (qI+pV)^{\lfloor n/2 \rfloor}\|_{TV} \\ &\quad + \frac{C(k)}{\sqrt{q}} n^{k+1} p^{3(k+1)} \|(V-I)^{*3(k+1)} * e^{npq(V-I)/6}\|_{TV} \\ &\leq \frac{C(k)p^{3(k+1)/2}}{n^{(k+1)/2} q^{(3k+4)/2}}. \end{aligned}$$

□

Lemma 8. *Let assumptions (4) hold. Then, for any $a > 0$, $k \in \mathbb{N}$, the following estimate holds*

$$\|(F-I)^{*k} * \exp(a(F-I))\|_{TV} \leq \left(\frac{k}{2a}\right)^k \delta^k(a/k) \leq \frac{C(k)\delta^k(a)}{a^k}, \quad (35)$$

where $\delta(a)$ is defined by (5).

Proof. Since F is symmetric, we can write

$$F-I = \sum_{\vec{x}} p_{\vec{x}} (I_{\vec{x}} - I) = \sum_{\vec{x}} p_{\vec{x}} (I_{-\vec{x}} - I) = \frac{1}{2} \sum_{\vec{x}} p_{\vec{x}} (I_{\vec{x}} + I_{-\vec{x}} - 2I). \quad (36)$$

Total variation of CP distribution equals 1. Therefore,

$$\exp(a(F-I)) = \exp(p_{\vec{x}} (I_{\vec{x}} + I_{-\vec{x}} - 2I)) * \Theta,$$

for any \vec{x} from F support and some $\Theta = \Theta(F)$ such that $\|\Theta\|_{TV} \leq 1$.

Therefore, applying (27) we get

$$\|(F-I) * \exp(a(F-I))\|_{TV}$$

$$\begin{aligned}
&\leq \frac{1}{2} \sum_{\bar{x}} p_{\bar{x}} \|(I_{\bar{x}} + I_{-\bar{x}} - 2I) * \exp(p_{\bar{x}}(I_{\bar{x}} + I_{-\bar{x}} - 2I))\|_{TV} \\
&\leq \frac{1}{2} \sum_{\bar{x}} p_{\bar{x}} \min\left(4, \frac{1}{ap_{\bar{x}}}\right) = \frac{1}{2a} \delta(a).
\end{aligned}$$

Estimate for $k > 1$ follows from (25). \square

Lemma 9. *Let assumptions (14)–(17) hold. Then, for any $a > 0$, $k \in \mathbb{N}$, the following estimate holds*

$$\|(F - I)^{*k} * \exp(a(F - I))\|_{TV} \leq \frac{(3.6k)^k}{e^k a^k} \left(\sum_{m=1}^K \sqrt{1 + \sigma_m} \right)^k. \quad (37)$$

Proof. Observe that $e^{a(F-I)}$ can be expressed as convolution of two CP distributions

$$e^{a(F-I)} = \exp(ap_m(F_m - I)) * \Theta, \quad \|\Theta\|_{TV} = \left\| \exp\left(a \sum_{j \neq m} p_j(F_j - I)\right) \right\|_{TV} = 1,$$

where the last equality follows from the fact that total variation norm of any distribution equals 1. Noting that estimates for convolutions of F_m can be treated as estimates for one-dimensional distributions (see (18)) and applying Lemma 4 we get

$$\begin{aligned}
\|(F - I) * \exp(a(F - I))\|_{TV} &\leq \sum_{m=1}^K p_m \|(F_m - I) * \exp(ap_m(F_m - I))\|_{TV} \\
&\leq \frac{3.6}{ae} \sum_{m=1}^K \sqrt{1 + \sigma_m}.
\end{aligned}$$

For the proof, when $k > 1$, we use (25). \square

Lemma 10. *Let $F = qI + pV \in \mathcal{F}_d$, $q \in (0, 1)$, and let D be defined by (8). Then for any $n \in \mathbb{N}$ the following equalities hold*

$$D^{*n} - e^{n(F-I)} = \frac{3.5n}{2\sqrt{q}} (F-I)^{*2} * e^{nq(F-I)/2} * \Theta, \quad (38)$$

$$D^{*n} - e^{n(F-I)} = -\frac{n}{2} e^{n(F-I)} * (F-I)^{*2} + \frac{3.5n^2}{4\sqrt{q}} e^{nq(F-I)/2} * (F-I)^{*4} * \Theta. \quad (39)$$

Proof. Observe that $F - I = p(V - I)$. Then applying (23) and (29) we obtain

$$\begin{aligned}
D^{*n} - e^{n(F-I)} &= e^{n(F-I)} * \left(e^{-n(F-I)^{*2}/2} - I \right) \\
&= -\frac{n}{2} e^{n(F-I)} * (F - I)^{*2} * \int_0^1 e^{-n\tau(F-I)^{*2}/2} d\tau \\
&= \frac{n}{2} e^{nq(F-I)/2} * (F - I)^{*2} * \int_0^1 (-1) e^{n\tau(1+p)(V-I)/2 - \tau n p^2 (V-I)^{*2}/2} d\tau
\end{aligned}$$

$$= \frac{3.5n}{2\sqrt{q}} e^{nq(F-I)/2} * (F-I)^{*2} * \Theta,$$

which proves (38). The proof of (39) is similar:

$$\begin{aligned} D^{*n} - e^{n(F-I)} &= -\frac{n}{2} e^{n(F-I)} * (F-I)^{*2} + e^{n(F-I)} * \left(e^{-n(F-I)^{*2}/2} - I + n(F-I)^{*2}/2 \right) \\ &= -\frac{n}{2} e^{n(F-I)} * (F-I)^{*2} + \frac{3.5n^2}{4\sqrt{q}} e^{nq(F-I)/2} * (F-I)^{*4} * \Theta. \end{aligned}$$

□

For the proof of Proposition 1 we need the following lemma.

Lemma 11. *Let $\vec{x} \in \mathbb{R}^d$, $\lambda > 0$. Then*

$$\begin{aligned} \left| \left\| (I_{-\vec{x}} + I_{\vec{x}} - 2I) * \exp(\lambda(I_{-\vec{x}} + I_{\vec{x}} - 2I)) \right\| - \frac{1}{\lambda} \sqrt{\frac{2}{\pi e}} \right| &\leq \frac{C}{\lambda\sqrt{\lambda}}, \\ \left| \left\| (I_{-\vec{x}} + I_{\vec{x}} - 2I)^{*2} * \exp(\lambda(I_{-\vec{x}} + I_{\vec{x}} - 2I)) \right\| - \frac{C^*}{\lambda^2} \right| &\leq \frac{C}{\lambda^2\sqrt{\lambda}}. \end{aligned}$$

Here

$$C^* = e^{-3/2} \sqrt{\frac{3}{\pi}} \left(e^{\sqrt{3/2}} \sqrt{3 - \sqrt{6}} + e^{-\sqrt{3/2}} \sqrt{3 + \sqrt{6}} \right).$$

For $d = 1$ both estimates were proved in [7], Lemma 4.7. In view of (26), the same proof holds for $d > 1$.

5 Proofs

Proofs of Theorem 1 and 7. We recall that total variation distance is half of the total variation norm. From Lemma 7 it follows that

$$\begin{aligned} \left\| F^{*n} - e^{n(F-I)} \right\|_{TV} &\leq \frac{C(1-q)^{9/2}}{n\sqrt{n}q^5} + \left\| D^{*n} - e^{n(F-I)} \right\|_{TV} \\ &+ \sum_{m=1}^2 \binom{n}{m} \left\| D^{*(n-m)} * (F-D)^{*m} \right\|_{TV}. \end{aligned} \quad (40)$$

Taking into account (32), (33) and (28) and observing that $\|F - I\|_{TV} \leq 2$ we obtain

$$\begin{aligned} \sum_{m=1}^2 \binom{n}{m} \left\| D^{*(n-m)} * (F-D)^{*m} \right\|_{TV} &\leq C \sum_{m=1}^2 n^m \|D^{-m}\|_{TV} \|D^{*n} * (F-I)^{*3m}\|_{TV} \\ &\leq \frac{C}{\sqrt{q}} \sum_{m=1}^2 n^m \left\| e^{nq(F-I)/2} * (F-I)^{*3m} \right\|_{TV} \leq \frac{C\sqrt{n}}{\sqrt{q}} \left\| e^{nq(F-I)/4} * (F-I)^{*2} \right\|_{TV} \\ &\times \sum_{m=1}^2 n^{m-1/2} \left\| e^{nq(F-I)/4} * (F-I)^{*(2m-1)} \right\|_{TV} \|F - I\|_{TV}^{m-1} \end{aligned}$$

$$\leq \frac{C\sqrt{n}}{q^2} \left\| e^{nq(F-I)/4} * (F-I)^{*2} \right\|_{TV}.$$

Applying (35) we get

$$\sum_{m=1}^2 \binom{n}{m} \left\| D^{*(n-m)} * (F-D)^{*m} \right\|_{TV} \leq \frac{C\delta^2(nq/4)}{n\sqrt{n}q^4} \leq \frac{C\delta^2(n/2)}{n\sqrt{n}q^4}. \quad (41)$$

Applying (37) we get

$$\sum_{m=1}^2 \binom{n}{m} \left\| D^{*(n-m)} * (F-D)^{*m} \right\|_{TV} \leq \frac{C}{n\sqrt{n}q^4} \left(\sum_{m=1}^K \sqrt{1+\sigma_m} \right)^2. \quad (42)$$

From (39) and (28) it follows that

$$\begin{aligned} \|D^{*n} - e^{n(F-I)}\|_{TV} &\leq \frac{n}{2} \left\| (F-I)^{*2} * e^{n(F-I)} \right\|_{TV} + \frac{3.5n^2}{4\sqrt{q}} \left\| (F-I)^{*4} * e^{nq(F-I)/2} \right\|_{TV} \\ &\leq \frac{n}{2} \left\| (F-I)^{*2} * e^{n(F-I)} \right\|_{TV} \\ &\quad + \frac{3.5n^2}{4\sqrt{q}} \left\| (F-I)^{*3} * e^{nq(F-I)/4} \right\|_{TV} \left\| (F-I) * e^{nq(F-I)/4} \right\|_{TV} \\ &\leq \frac{n}{2} \left\| (F-I)^{*2} * e^{n(F-I)} \right\|_{TV} + \frac{Cn\sqrt{n}}{q} \left\| (F-I)^{*3} * e^{nq(F-I)/4} \right\|_{TV}. \end{aligned}$$

Applying (35) we get

$$\|D^{*n} - e^{n(F-I)}\|_{TV} \leq \frac{\delta^2(n/2)}{2n} + \frac{C\delta^3(nq/4)}{n\sqrt{n}q^4} \leq \frac{\delta^2(n/2)}{2n} + \frac{C\delta^2(n/2)\delta(nq)}{n\sqrt{n}q^4}. \quad (43)$$

Applying (37) we get

$$\|D^{*n} - e^{n(F-I)}\|_{TV} \leq \frac{25.92}{e^2 n} \left(\sum_{m=1}^K \sqrt{1+\sigma_m} \right)^2 + \frac{C}{n\sqrt{n}q^4} \left(\sum_{m=1}^K \sqrt{1+\sigma_m} \right)^3. \quad (44)$$

Substituting (41) and (43) into (40) we complete the proof of (6). Substituting (42) and (44) into (40) we complete the proof of (19). \square

Proofs of Theorem 1* and Theorem 7*. Applying Lemma 7, (38), (32), (33) we get

$$\begin{aligned} \|F^{*n} - e^{n(F-I)}\|_{TV} &\leq \|M_1\|_{TV} + \|D^{*n} - e^{n(F-I)}\|_{TV} + n \left\| D^{*(n-1)} * (F-D) \right\|_{TV} \\ &\leq \frac{C(1-q)^3}{nq^{7/2}} + \frac{Cn}{\sqrt{q}} \left\| (F-I)^{*2} * e^{nq(F-I)/4} \right\|_{TV} \left\| e^{nq(F-I)/4} \right\|_{TV} \\ &\quad + \frac{Cn}{\sqrt{q}} \|D^{*(-1)}\|_{TV} \left\| (F-I)^{*2} * e^{nq(F-I)/4} \right\|_{TV} \|F-I\|_{TV} \left\| e^{nq(F-I)/4} \right\|_{TV} \\ &\leq \frac{C(1-q)^3}{nq^{7/2}} + \frac{Cn}{\sqrt{q}} \left\| (F-I)^{*2} * e^{nq(F-I)/4} \right\|_{TV}. \end{aligned}$$

It remains to apply (35) or (37). \square

Proof of Theorem 2. From Lemma 7 it follows that

$$\|F^{*n} - D^{*n}\|_{TV} \leq \frac{C(1-q)^{15/2}}{q^8 n^2 \sqrt{n}} + \sum_{m=1}^4 \binom{n}{m} \|D^{*(n-m)} * (F-D)^{*m}\|_{TV}. \quad (45)$$

Therefore, applying (32), (33), (27) and (35) we prove that

$$\begin{aligned} \sum_{m=2}^4 \binom{n}{m} \|D^{*(n-m)} * (F-D)^{*m}\|_{TV} &\leq \sum_{m=2}^4 \binom{n}{m} \|D^{*n} * (F-D)^{*m}\|_{TV} e^{4m} \\ &\leq \frac{C}{\sqrt{q}} \sum_{m=2}^4 n^m \|e^{nq(F-I)/2} * (F-I)^{*3m}\|_{TV} \leq \frac{C}{\sqrt{q}} \|e^{nq(F-I)/4} * (F-I)^{*3}\|_{TV} \\ &\times \sum_{m=2}^4 n^m \|e^{nq(F-I)/4} * (F-I)^{*2m-1}\|_{TV} \|F-I\|_{TV}^{m-2} \\ &\leq \frac{C}{\sqrt{q}} \|e^{nq(F-I)/12} * (F-I)\|_{TV}^3 \sum_{m=2}^4 \frac{n^m 2^{m-2}}{(nq)^{m-1/2}} \leq \frac{C\delta^3(nq/12)}{q^7 n^2 \sqrt{n}} \\ &\leq \frac{C\delta^3(nq/6)}{q^7 n^2 \sqrt{n}}. \end{aligned} \quad (46)$$

Similarly applying (33), (31), (32), (28) and (35) we get

$$\begin{aligned} n\|D^{*(n-1)} * (F-D)\|_{TV} &\leq \frac{n}{3} \|D^{*(n-1)} * (F-I)^{*3}\|_{TV} + Cn\|D^{*(n-1)} * (F-I)^{*4}\|_{TV} \\ &\leq \frac{n}{3} \|D^{*n} * (F-I)^{*3}\|_{TV} + \frac{n}{3} \|D^{*(-1)}\|_{TV} \|D^{*n} * (I-D) * (F-I)^{*3}\|_{TV} \\ &+ Cn\|D^{*(-1)}\|_{TV} \|D^{*n} * (F-I)^{*4}\|_{TV} \\ &\leq \frac{3.5n}{3\sqrt{q}} \|e^{nq(F-I)/2} * (F-I)^{*3}\|_{TV} + \frac{Cn}{\sqrt{q}} \|e^{nq(F-I)/2} * (F-I)^{*4}\|_{TV} \\ &\leq \frac{31.5\delta^3(nq/6)}{q^{7/2}n^2} + \frac{Cn}{\sqrt{q}} \|e^{nq(F-I)/4} * (F-I)\|_{TV} \|e^{nq(F-I)/4} * (F-I)^{*3}\|_{TV} \\ &\leq \frac{31.5\delta^3(nq/6)}{q^{7/2}n^2} + \frac{C\sqrt{n}}{q} \|e^{nq(F-I)/4} * (F-I)^{*3}\|_{TV} \\ &\leq \frac{31.5\delta^3(nq/6)}{q^{7/2}n^2} + \frac{C\delta^3(nq)}{q^4 n^2 \sqrt{n}}. \end{aligned} \quad (47)$$

Collecting estimates (45), (46) and (47) we complete the proof of Theorem. \square

Proofs of Theorems 3 and 9. From Lemma 7 it follows that

$$\begin{aligned} \|F^{*n} - e^{n(F-I)} * \left(I - \frac{n}{2}(F-I)^{*2}\right)\|_{TV} \\ \leq \frac{C(1-q)^6}{n^2 q^{13/2}} + \sum_{m=1}^3 \binom{n}{m} \|D^{*(n-m)} * (F-D)^{*m}\|_{TV} \end{aligned}$$

$$+ \left\| D^{*n} - e^{n(F-I)} * \left(I - n(F-I)^{*2}/2 \right) \right\|_{TV} \quad (48)$$

Taking into account (32), (33) and (28) we obtain

$$\begin{aligned} \sum_{m=1}^3 \binom{n}{m} \left\| D^{*(n-m)} * (F-D)^{*m} \right\|_{TV} &\leq C \sum_{m=1}^3 \binom{n}{m} \|D^{*(-m)}\|_{TV} \|D^{*n} * (F-I)^{*3m}\|_{TV} \\ &\leq \frac{C}{\sqrt{q}} \sum_{m=1}^3 \binom{n}{m} e^{4m} \left\| e^{nq(F-I)/2} * (F-I)^{*3m} \right\|_{TV} \\ &\leq \frac{Cn}{\sqrt{q}} \left\| e^{nq(F-I)/4} * (F-I)^{*3} \right\|_{TV} \\ &\times \sum_{m=1}^3 n^{m-1} \left\| e^{nq(F-I)/4} * (F-I)^{*2(m-1)} \right\|_{TV} \|F-I\|_{TV}^{m-1} \\ &\leq \frac{Cn}{\sqrt{q}} \left\| e^{nq(F-I)/4} * (F-I)^{*3} \right\|_{TV} \sum_{m=2}^3 \frac{n^{m-1} 2^{m-1}}{n^{m-1} q^{m-1}} \\ &\leq \frac{Cn}{q^{5/2}} \left\| e^{nq(F-I)/4} * (F-I)^{*3} \right\|_{TV}. \end{aligned} \quad (49)$$

From (39) it follows that

$$\left\| D^{*n} - e^{n(F-I)} * \left(I - \frac{n}{2} (F-I)^{*2} \right) \right\|_{TV} \leq \frac{Cn^2}{\sqrt{q}} \left\| e^{nq(F-I)/2} * (F-I)^{*4} \right\|_{TV}. \quad (50)$$

To proof (10) (respectively (22)) it suffices in (49)–(50) to use (35) (respectively (37)) and substitute resulting estimates into (48). \square

Proofs of Theorems 4 and 10. By the triangle inequality

$$\begin{aligned} \left\| F^{*n} - F^{*(n+1)} \right\|_{TV} &\leq \left\| F^{*n} - e^{n(F-I)} \right\|_{TV} + \left\| e^{n(F-I)} - e^{(n+1)(F-I)} \right\|_{TV} \\ &+ \left\| F^{*(n+1)} - e^{(n+1)(F-I)} \right\|_{TV}. \end{aligned}$$

From definition of exponential measure it follows that

$$\left\| e^{n(F-I)} - e^{(n+1)(F-I)} \right\|_{TV} = \left\| e^{n(F-I)} * (I - e^{F-I}) \right\|_{TV} \leq C \left\| e^{n(F-I)} * (F-I) \right\|_{TV}.$$

It remains to apply (7) and (35) or (20) and (37). \square

Proof of Theorem 5. Without loss of generality let $a < b$. From definition of exponential measure (23), the fact that total variation of any distribution equals 1 and Lemma 8 we obtain

$$\begin{aligned} \left\| e^{b(F-I)} - e^{a(F-I)} \right\|_{TV} &= \left\| e^{a(F-I)} * \left(e^{(b-a)(F-I)} - I \right) \right\|_{TV} \\ &= \left\| (b-a) e^{a(F-I)} * (F-I) * \int_0^1 e^{\tau(b-a)(F-I)} d\tau \right\|_{TV} \end{aligned}$$

$$\begin{aligned}
&\leq (b-a) \left\| e^{a(F-I)} * (F-I) \right\|_{TV} \int_0^1 \left\| e^{\tau(b-a)(F-I)} \right\|_{TV} d\tau \\
&\leq (b-a) \left\| e^{a(F-I)} * (F-I) \right\|_{TV} \leq \frac{2(b-a)\delta(a)}{ae} \\
&\leq \frac{2(b-a)(2N+1)}{ae}.
\end{aligned} \tag{51}$$

If $b \leq a(1 + e/(2N+1))$, then

$$\frac{2(b-a)(2N+1)}{ae} \leq \frac{2(b-a)}{b} \left(1 + \frac{2N+1}{e}\right)$$

and theorem's statement follows from (51). If $b > a(1 + e/(2N+1))$, then directly

$$\left\| e^{b(F-I)} - e^{a(F-I)} \right\|_{TV} \leq \left\| e^{b(F-I)} \right\|_{TV} + \left\| e^{a(F-I)} \right\|_{TV} = 2 \leq \frac{2(b-a)}{b} \left(1 + \frac{2N+1}{e}\right).$$

Recall that total variation distance is half of the total variation norm. \square

Proof of Theorem 6. Applying Lemma 7, (32), (28), (33) and (35) we obtain

$$\begin{aligned}
\|M_s\|_{TV} &\leq \|M_{4s+3}\|_{TV} + \sum_{m=s+1}^{4s+3} \binom{n}{m} \left\| D^{*(n-m)} * (F-D)^{*m} \right\|_{TV} \\
&\leq \frac{C(s)(1-q)^{6(s+1)}}{n^{2(s+1)}q^{6s+13/2}} + C \sum_{m=s+1}^{4s+3} n^m \|D^{*(-m)}\|_{TV} \|D^{*n} * (F-I)^{*3m}\|_{TV} \\
&\leq \frac{C(s)(1-q)^{6(s+1)}}{n^{2(s+1)}q^{6s+13/2}} + \frac{C(s)}{\sqrt{q}} \sum_{m=s+1}^{4s+3} n^m \left\| e^{nq(F-I)/2} * (F-I)^{*3m} \right\|_{TV} \\
&\leq \frac{C(s)(1-q)^{6(s+1)}}{n^{2(s+1)}q^{6s+13/2}} + \frac{C(s)n^{s+1}}{\sqrt{q}} \left\| e^{nq(F-I)/4} * (F-I)^{*3(s+1)} \right\|_{TV} \\
&\times \left(1 + \sum_{m=1}^{3s+2} n^m \left\| e^{nq(F-I)/4} * (F-I)^{*2m} \right\|_{TV} \|F-I\|_{TV}^m \right) \\
&\leq \frac{C(s)(1-q)^{6(s+1)}}{n^{2(s+1)}q^{6s+13/2}} + \frac{C(s)\delta^{3(s+1)}(nq)}{q^{6s+11/2}n^{2(s+1)}}.
\end{aligned}$$

\square

Proof of Theorem 8. Applying Lemma 7, (32), (33), (28) and (37) we get

$$\begin{aligned}
\|F^{*n} - D^{*n}\|_{TV} &\leq \|M_3\|_{TV} + C \sum_{m=1}^3 \binom{n}{m} \|D^{*(-m)}\|_{TV} \|D^{*n} * (F-I)^{*3m}\|_{TV} \\
&\leq \frac{C(1-q)^6}{n^2q^{13/2}} + \frac{C}{\sqrt{q}} \sum_{m=1}^3 e^{4m} n^m \left\| e^{nq(F-I)/2} * (F-I)^{*3m} \right\|_{TV} \\
&\leq \frac{C(1-q)^6}{n^2q^{13/2}} + \frac{Cn}{\sqrt{q}} \left\| e^{nq(F-I)/4} * (F-I)^{*3} \right\|_{TV}
\end{aligned}$$

$$\begin{aligned}
& \times \sum_{m=1}^3 n^{m-1} \left\| e^{nq(F-I)/4} * (F-I)^{*2(m-1)} \right\|_{TV} 2^{m-1} \\
& \leq \frac{C(1-q)^6}{n^2 q^{13/2}} + \frac{C}{n^2 q^{7/2}} \left(\sum_{m=1}^K \sqrt{1+\sigma_m} \right)^3 \sum_{m=2}^3 \frac{n^{m-1} 2^{m-1}}{n^{m-1} q^{m-1}} \\
& \leq \frac{C}{n^2 q^{13/2}} \left(\sum_{m=1}^K \sqrt{1+\sigma_m} \right)^3.
\end{aligned}$$

□

Proof of Theorem 11. Without loss of generality let $a < b$. Repeating the proof of (51) but using (37) we prove that

$$\left\| e^{b(F-I)} - e^{a(F-I)} \right\|_{TV} \leq (b-a) \left\| e^{a(F-I)} * (F-I) \right\|_{TV} \leq \frac{3.6(b-a)}{ae} \sum_{m=1}^K \sqrt{1+\sigma_m}.$$

If $b \leq 2.5a$, then

$$\frac{3.6(b-a)}{ae} \sum_{m=1}^K \sqrt{1+\sigma_m} \leq \frac{3.4(b-a)}{b} \sum_{m=1}^K \sqrt{1+\sigma_m}.$$

and theorem's statement follows from analogue of (51). If $b > 2.5a$, then directly

$$\left\| e^{b(F-I)} - e^{a(F-I)} \right\|_{TV} \leq 2 \leq \frac{3.4(b-a)}{b} \leq \frac{3.4(b-a)}{b} \sum_{m=1}^K \sqrt{1+\sigma_m}.$$

□

Proof of Proposition 1. All characteristics of F are constants. Therefore, we can write C instead of $C(F)$. From the proof of Theorem 1 we see that

$$n \| F^{*n} - \exp(n(F-I)) \| \leq \frac{n^2}{2} \left\| (F-I)^{*2} * e^{n(F-I)} \right\|_{TV} + \frac{C}{\sqrt{n}}.$$

By assumptions

$$\begin{aligned}
F - I &= \sum_{j=1}^N p_{\bar{x}_j} \left(I_{-\bar{x}_j} + I_{\bar{x}_j} - 2I \right), \quad (F-I)^{*2} = \sum_{j=1}^N p_{\bar{x}_j}^2 \left(I_{-\bar{x}_j} + I_{\bar{x}_j} - 2I \right)^{*2} \\
&+ \sum_{j \neq k}^N p_{\bar{x}_j} \left(I_{-\bar{x}_j} + I_{\bar{x}_j} - 2I \right) p_{\bar{x}_k} \left(I_{-\bar{x}_k} + I_{\bar{x}_k} - 2I \right) \\
\exp(n(F-I)) &= \exp\left(np_{\bar{x}_j} \left(I_{-\bar{x}_j} + I_{\bar{x}_j} - 2I \right)\right) * \exp\left(np_{\bar{x}_k} \left(I_{-\bar{x}_k} + I_{\bar{x}_k} - 2I \right)\right) * \Theta \\
&= \exp(n(F-I)) = \exp\left(np_{\bar{x}_j} \left(I_{-\bar{x}_j} + I_{\bar{x}_j} - 2I \right)\right) * \Theta.
\end{aligned}$$

Therefore, taking into account Lemma 11, we get

$$\begin{aligned} & \frac{n^2}{2} \left\| (F - I)^{*2} * e^{n(F-I)} \right\|_{TV} \\ & \leq \frac{n^2}{2} \sum_{j=1}^N p_{\bar{x}_j}^2 \left\| (I_{-\bar{x}_j} + I_{\bar{x}_j} - 2I)^{*2} * \exp \left(np_{\bar{x}_j} (I_{-\bar{x}_j} + I_{\bar{x}_j} - 2I) \right) \right\|_{TV} \\ & + \frac{n^2}{2} \sum_{j \neq k}^N p_{\bar{x}_j} \left\| (I_{-\bar{x}_j} + I_{\bar{x}_j} - 2I) * \exp \left(np_{\bar{x}_j} (I_{-\bar{x}_j} + I_{\bar{x}_j} - 2I) \right) \right\|_{TV} \\ & \times p_{\bar{x}_k} \left\| (I_{-\bar{x}_k} + I_{\bar{x}_k} - 2I) * \exp \left(np_{\bar{x}_k} (I_{-\bar{x}_k} + I_{\bar{x}_k} - 2I) \right) \right\|_{TV} \\ & \leq \frac{C^* N}{2} + \frac{N(N-1)}{\pi e} + \frac{C}{\sqrt{n}}. \end{aligned}$$

To complete the proof recall that the total variation metric is half of the total variation norm. □

Remarks on simulations. R-studio, library(cubature) and formulas of inversion were used. Example of the program is given below

```

1 >n<-10; p1<-0.10; p2<-0.10; q1<-1-2*p1; q2<-1-2*p2; S<-0; Z<-0; U<-0;
  for (m in -n:n){integrandz1<- function(x) (q1+2*p1*cos(x[1]))^n*cos(m*x[1])/(2*pi); z1= hcubature(integrandz1, lowerLimit=-pi,
  upperLimit=pi); integranda1<- function(x) exp(2*n*p1*(cos(x[1])
  -1))*cos(x[1]*m)/(2*pi); a1= hcubature(integranda1,lowerLimit=-pi,
  upperLimit=pi);
2 for (k in -n:n){integrandz2<- function(x) (q2+2*p2*cos(x[1]))^n*cos(k
  *x[1])/(2*pi); z2= hcubature(integrandz2, lowerLimit=-pi,
  upperLimit=pi); integranda2<- function(x) exp(2*n*p2*(cos(x[1])-1)
  )*cos(x[1]*k)/(2*pi); a2= hcubature(integranda2, lowerLimit=-pi,
  upperLimit=pi); S=S+abs(z1$integral*z2$integral-a1$integral*a2$
  integral); Z=Z+ a1$integral*a2$integral; U<-U+z1$integral*z2$
  integral }; print(U); print(S); print(Z); var<-S+1-Z; dtv<-var/
  2; print(dtv)
3

```

Acknowledgment

We are grateful to the referees for their valuable remarks that allowed to improve the initial version of the paper.

References

[1] Barbour, A.D., Holst, L., Janson, S.: Poisson Approximation. Clarendon Press, Oxford (1992). [MR1163825](#)

[2] Barbour, A.D., Chryssaphinou, O., Vaggelatou, E.: Applications of compound Poisson approximation. In: Charalambides, C.A., Koutras, M.V., Balakrishnan, N. (eds.) Probability and Statistical Models with Applications, pp. 41–62. Chapman & Hall (2001). [MR1865030](#). <https://doi.org/10.1214/aoap/1015345355>

- [3] Barman, K., Upadhye, N.S., Vellaisamy, P.: Approximations related to tempered stable distributions. *Mod. Stoch. Theory Appl.* **12**(3), 325–346 (2025). [MR4917904](#). <https://doi.org/10.15559/25-vmsta275>
- [4] Barbour, A.D., Luczak, M.J., Xia, A.: Multivariate approximation in total variation II: discrete normal approximation. *Ann. Probab.* **46**, 1405–1440 (2018). [MR3785591](#). <https://doi.org/10.1214/17-AOP1205>
- [5] Bergström, H.: On asymptotic expansion of probability functions. *Skand. Aktuarietidskr.* **1**, 1–34 (1951). [MR0043403](#). <https://doi.org/10.1080/03461238.1951.10432122>
- [6] Bobkov, S.G., Götze, F.: Berry–Esseen bounds in the local limit theorems. *Lith. Math. J.* **65**, 50–66 (2025). [MR4885697](#). <https://doi.org/10.1007/s10986-025-09659-1>
- [7] Čekanavičius, V., Roos, B.: Compound binomial approximations. *Ann. Inst. Stat. Math.* **58**, 187–210 (2006). [MR2281210](#). <https://doi.org/10.1007/s10463-005-0018-4>
- [8] Čekanavičius, V., Novak, S.: Compound Poisson approximation. Chapman & Hall/CRC Press, Boca Raton, London, New York (2024)
- [9] Genest, C., Marceau, E., Mesfioui, M.: Compound Poisson approximations for individual models with dependent risks. *Insur. Math. Econ.* **32**(1), 73–91 (2003). [MR3600678](#). <https://doi.org/10.1016/j.insmatheco.2016.10.012>
- [10] Götze, F., Yu, Z.A.: On alternative approximating distributions in the multivariate version of Kolmogorov’s second uniform limit theorem. *Teor. Veroâtn. Primen.* **61**(1), 3–22 (2022). Transl.: *Theory Probab. Appl.*, 67, 1, 1–16. [MR4466409](#). <https://doi.org/10.1137/s0040585x97t99071x>
- [11] Hipp, C.: Improved approximations for the aggregate claims distribution in the individual model. *ASTIN Bull.* **16**(2), 89–100 (1986). <https://doi.org/10.2143/AST.16.2.2015001>
- [12] Jokubauskienė, S., Čekanavičius, V.: Multivariate Hipp-type compound Poisson approximations for lattice distributions I. *Lith. Math. J.* **65**, 337–356 (2025). [MR4953667](#). <https://doi.org/10.1007/s10986-025-09686-y>
- [13] Kruopis, J.: Precision of approximations of the generalized Binomial distribution by convolutions of Poisson measures. *Litovsk. Mat. Sb.* **26**(1), 53–69 (1986). (Russian). Transl.: *Lith. Math. J.*, 26(1), 37–49. [MR0847204](#)
- [14] Kruopis, J., Čekanavičius, V.: Compound Poisson approximations for symmetric vectors. *J. Multivar. Anal.* **123**, 30–42 (2014). [MR3130419](#). <https://doi.org/10.1016/j.jmva.2013.08.017>
- [15] Lishamol, T., Veena, G.: A retrospective study of Skellam and related distributions. *Austrian J. Stat.* **51**, 102–111 (2022). <https://doi.org/10.17713/ajs.v51i1.1224>
- [16] Lu, D., Chen, Y., Wang, L.: Asymptotics for the joint tail probability of bidimensional randomly weighted sums of dependent and heavy-tailed random variables. *Lith. Math. J.* **66**, 79–92 (2026). [MR5022834](#). <https://doi.org/10.1007/s10986-026-09709-2>
- [17] Novak, S.Y.: Extreme value methods with applications to finance. Chapman & Hall/CRC Press, London (2011). [MR2933280](#)
- [18] Roos, B.: Binomial approximation to the Poisson binomial distribution: the Krawtchouk expansion. *Teor. Veroâtn. Primen.* **45**(2), 328–344 (2000). Reprinted in: *Theory Probab. Appl.*, 45(2), 258–272. [MR1967760](#). <https://doi.org/10.4213/tvp466>
- [19] Roos, B.: Multinomial and Krawtchouk approximations to the generalized multinomial distribution. *Teor. Veroâtn. Primen.* **46**(1), 117–133 (2001). Reprinted in: *Theory Probab. Appl.* 46(1), 103–117. [MR1968708](#). <https://doi.org/10.4213/tvp3954>
- [20] Roos, B.: Kerstan’s method in the multivariate Poisson approximation: an expansion in the exponent. *Teor. Veroâtn. Primen.* **43**(2), 397–402 (2002). Reprinted in: *Theory Probab. Appl.*, 47(2), 358–363. [MR2003208](#). <https://doi.org/10.4213/tvp3672>

- [21] Roos, B.: On variational bounds in the compound Poisson approximation of the individual risk model. *Insur. Math. Econ.* **40**(3), 403–414 (2007). [MR2310979](#). <https://doi.org/10.1016/j.insmatheco.2006.06.003>
- [22] Roos, B.: Refined total variation bounds in the multivariate and compound Poisson approximation. *ALEA Lat. Am. J. Probab. Math. Stat.* **14**, 337–360 (2017). [MR3647296](#). <https://doi.org/10.30757/alea.v14-19>
- [23] Sengar, A.S., Upadhye, N.S.: Subordinated compound Poisson processes of order k . *Mod. Stoch. Theory Appl.* **7**(4), 395–413 (2020). [MR4195643](#). <https://doi.org/10.15559/20-vmsta165>
- [24] Sundt, B., Vernic, R.: *Recursions for Convolutions and Compound Distributions with Insurance Applications*. EAA Lecture Notes. Springer, Dordrecht, Heidelberg, London, New York (2009). [MR2502743](#). <https://doi.org/10.1007/978-3-540-92900-0>
- [25] Vellaisamy, P., Chaudhuri, B.: Poisson and compound Poisson approximations for random sums or random variables. *J. Appl. Probab.* **33**, 127–137 (1996). [MR1371960](#). <https://doi.org/10.2307/3215270>
- [26] Yu, Z.A.: Estimates for the closeness of successive convolutions of multidimensional symmetric distributions. *Probab. Theory Relat. Fields* **79**, 175–200 (1988). [MR0958287](#). <https://doi.org/10.1007/BF00320918>
- [27] Yu, Z.A.: Approximation of convolutions of multi-dimensional symmetric distributions by accompanying laws. *J. Sov. Math.* **177**(1), 55–72 (1989). (Russian). *Transl.*, 61, 1859–1872, 1992. [MR1053124](#). <https://doi.org/10.1007/BF01362793>
- [28] Yu, Z.A.: On Proximity of Distributions of Successive Sums with Respect to the Prokhorov Distance. *Teor. Veroyatn. Primen.* **69**(2), 272–284 (2024). (Russian). *Transl.: Theory Prob. Appl.* **69**(2), 217–226. [MR4912064](#). <https://doi.org/10.4213/typ5690>